

# MANZIL

## FOR JEE ASPIRANTS



**Physics**

**ATDB.uno**

**Circular motion**

**In One Shot** **By – Rajwant Singh (RJ)**



# Topics *to be covered*

- 1 Circular kinematics
- 2 Questions
- 3 Circular dynamics
- 4 Questions

ATDB.uno



— FOR NOTES & DPP CHECK DESCRIPTION —

# PWW MANANZIL IIT

## TELEGRAM CHANNEL



ATDB.uno



@PWJEEWALLAH





# Circular Motion

"Plane"



$x \rightarrow$  displacement  $\Rightarrow \Theta$  "Angular displacement".

$v = \frac{dx}{dt} \Rightarrow$  velocity  $\Rightarrow \omega$  "Angular velocity".

$a = \frac{dv}{dt} \Rightarrow$  acceleration  $\Rightarrow \alpha$  "Angular acceleration".



# Angular Displacement



$\Rightarrow \theta = \text{Angle Rotated by particle.}$



$\Rightarrow \frac{\text{Arc length}}{\text{Radius}}$

$\Rightarrow \text{Radians} \Rightarrow [M^0 L^0 T^0]$

$\Rightarrow 180^\circ = \pi \text{ radians}$

$\Rightarrow$  direction by RHTR.  
 Right hand  $\rightarrow$  finger  $\rightarrow$  dir of Motion  
 Thumb  $\rightarrow$   $\theta$  dir.

on axis of Rotation.



ATDB.uno



# Angular Velocity





$\vec{\omega}$  = Angular Velocity  
 ⇒ Rate of change of Angular displacement with time.

$$\omega = \frac{d\theta}{dt}$$

unit =  $\frac{\text{rad}}{\text{s}}$

Dimension =  $[T^{-1}]$



ATDB.uno

⊙ Instantaneous angular velocity.

$\theta = 2t^2$  (Circular Motion)

$$\omega = \frac{d\theta}{dt} = \frac{d}{dt}(2t^2) = 4t$$

on axis of Rotation.

$\omega$  at any Instant.  $\omega$  at  $t = 1 \text{ sec}$ .

$$\omega_{t=1s} = 4 \text{ rad/s}$$

⊙ RHTR.

Right hand finger → dir of Motion.

Thumb →  $\omega$ .




# ⊙ Average angular velocity

$$\omega_{av} = \frac{\text{Total Angle travelled}}{\text{Total time}} = \frac{\theta_2 - \theta_1}{t_2 - t_1}$$

#

⊙ If particle Cover 1 Rotation with Constant Speed in time = T.

$$\omega = \frac{\text{Total Angle}}{\text{Total time}} = \frac{2\pi}{T}$$



$$\omega = \frac{2\pi}{T}$$
$$\omega = 2\pi \nu$$

$$\text{Unit} = \text{rps, rpm.}$$

$$\frac{1}{T} = \text{frequency} = \nu \Rightarrow$$

"No of Rotations per Second"

# 120 rpm

$$\text{rps} = \frac{\text{rpm}}{60}$$
$$\nu = \frac{120 \text{ rpm}}{60} = 2 \text{ rps}$$
$$\omega = 2\pi \nu = 4\pi \text{ rad/s.}$$



# Angular Acceleration



$\Rightarrow \alpha =$  Rate of change of angular velocity w.r.t time.

$$\alpha = \frac{d\omega}{dt}$$

$$\text{Unit} = \frac{\text{rad}}{\text{s}^2}$$

$$\text{Dimension} = [T^{-2}]$$

ATDB.uno

⊙ Instantaneous angular acceleration

$$\theta = 2t^3 + 3t^2 + 4$$

Find  $\alpha$  at  $t = 1 \text{ sec}$ .

$\rightarrow$  Speed up.

$$\# \omega = \frac{d\theta}{dt} = 6t^2 + 6t$$

$$\# \alpha = \frac{d\omega}{dt} = 12t + 6 \Rightarrow \alpha_{t=1} = 12 + 6 = 18 \frac{\text{rad}}{\text{s}^2}$$



Average Angular Acceleration


$$a_{rav} = \frac{\omega_f - \omega_i}{t_f - t_i}$$

⊙ Kinematics

"Retardation" ⇒ Rukna.

$a = \oplus$  or  $\ominus$  both ho sakti hai.

ATDB.uno



$v$  & acc are anti parallel.

Similar in Circular Kinematics.

$\vec{\omega}$  &  $\vec{\alpha}$  (axial vectors)

"Aise vectors jo axis of Rotation par lagte hai.

$\omega = \text{Constant}$

Ex  $\omega = 5 \frac{\text{rad}}{\text{s}}$


$$\alpha = \frac{d\omega}{dt} = 0$$

Neither Speed up  
Nor Slow down.



ATDB.uno

Speed up.



Slow down.




# # Important Relation.



Q)

There is a Relation between.

$v$  Tangential & Angular velocity.



ATDB.uno

$$\vec{v} = \vec{\omega} \times \vec{R}$$

$$v = \omega R \sin 90$$

$$v = R\omega$$


$$\vec{\omega} = i + j - k$$

$$\vec{R} = 2i - j + 3k$$

$$\vec{v} = \begin{vmatrix} i & j & k \\ 1 & 1 & -1 \\ 2 & -1 & 3 \end{vmatrix} = i(3 - (-1)) - j(3 - (-2)) + k(-1 - 2)$$



b) Relation between  $a_t$  &  $\alpha$ .



$\odot \vec{v} = \omega \times \vec{R}$

Rate of change.

$$\frac{d}{dt} \vec{v}_T = \frac{d}{dt} (\vec{\omega} \times \vec{R})$$

$$a_T = \alpha \times R$$

$$a = \alpha R \sin 90$$

$$a = R\alpha$$

ATDB.uno



# Kinematics of Circular Motion



$$\textcircled{1} \quad \omega = \frac{d\theta}{dt}$$

$$\omega_{av} = \frac{\theta_2 - \theta_1}{t_2 - t_1}$$

$$\textcircled{2} \quad \alpha_r = \frac{d\omega}{dt} = \frac{d\omega}{dt} \cdot \frac{d\theta}{d\theta} = \omega \frac{d\omega}{d\theta}$$

$$\alpha_{var} = \frac{\omega_2 - \omega_1}{t_2 - t_1}$$

$$\textcircled{3} \quad \alpha_r = \text{Constant.}$$


$$\omega_f = \omega_0 + \alpha_r t$$

$$\theta = \omega_0 t + \frac{1}{2} \alpha_r t^2$$


$$\omega_f^2 - \omega_0^2 = 2\alpha_r \theta$$

$$\theta_{n^{th}} = \omega_0 + \frac{\alpha_r}{2} (2n-1)$$

# Graphs.



Slope =  $\omega$ .



Slope =  $\alpha$


Area =  $\theta_2 - \theta_1$



Area =  $\omega_2 - \omega_1$



# Centripetal and Tangential Acceleration



Koi bhi particle agar CM mein hai toh  
ek acc towards Centre toh hogi (Must)

⇒ Jo acc towards Centre hai & direction  
change kar rahi hai ⇒ Centripetal Acceleration

$$a_c = a_{\text{centripetal}} = \frac{v^2}{R} = R\omega^2$$

Towards Centre.

direction change karane ka kaam.


here are two type of C.M.

UCM  $\Rightarrow$   $a_{net}$  is towards Centre.

uniform Circular Motion

$\omega = \text{Constant}$  Ex  $\omega = 5 \frac{\text{rad}}{\text{s}}$ ,  $R = 2\text{m}$

$|v_{\text{Tangential}}| = \text{Constant}$   $v = R\omega = 10 \text{ m/s}$



UCM

$a_c \neq 0$   
 $a_c = R\omega^2 = \frac{v^2}{R}$   
 $a_t = 0$


$a_{net} \neq 0$  Not toward Centre.

NUCM

(Non-uniform C.M.)

$\omega = \text{Vary}$   
 Ex:  $\omega = 2t^2$   
 OR  
 $v = \text{Vary}$


Speed up



$a_c \neq 0 = \frac{v^2}{R} = R\omega^2$

$v \rightarrow \text{Vary}$   $a_c \rightarrow \text{Vary}$

Slow down.



#  $a_{\text{tangential}} = R\alpha = \frac{dv}{dt}$



ATDB.uno



### QUESTION 01

If  $\theta$  depends on time  $t$  in following way

$$\theta = 2t^2 + 3 \text{ then } \begin{matrix} t=0 & \theta = 2 \cdot 0^2 + 3 = 3 \\ t=3 & \theta = 2 \cdot 9 + 3 = 21 \end{matrix}$$

(a) Find out  $\omega$  average upto 3 sec.

(b)  $\omega$  at 3 sec (Instant)

$$a) \omega_{avg} = \frac{\theta_2 - \theta_1}{t_2 - t_1} = \frac{21 - 3}{3 - 0} = \frac{18}{3} = 6 \frac{\text{rad}}{\text{s}}$$

$$b) \omega = \frac{d\theta}{dt} = 4t$$

at  $t=3$

$$\omega = 4 \times 3 = 12 \frac{\text{rad}}{\text{s}}$$

### QUESTION 02

A particle moves in a circular path of radius 1 m with an angular speed

$$\omega = 2t^2 + 1 \text{ rad/sec}$$

$t \rightarrow \text{Vary } \omega \rightarrow \text{Vary } v \rightarrow \text{Vary}$


Find the angle between total acceleration and normal acceleration at  $t = 1$  sec.

Centripetal Acc.

Speed up.

Sol.:-

$$R = 1 \text{ m}$$



$$a_c = \frac{v^2}{R} = R\omega^2$$

$$\text{at } t=1 \quad a_c = R\omega^2 = 1 \times (3)^2 = 9$$

$$a_t = R\alpha = 1 \times 4 = 4 \text{ m/s}^2$$

$$\alpha = \frac{d\omega}{dt} = 4t \quad \text{at } t=1 \quad \alpha = 4$$



### QUESTION 03

If a particle moves in a circle describing equal angles in equal times, its velocity vector

- A** ~~Remains constant~~
- B** ~~Changes in magnitude~~
- C** Change in direction
- D** ~~Changes both in magnitude and direction~~

$$\omega = \frac{d\theta}{dt}$$

$$\omega = \text{Constant}$$

$$|\mathbf{v}| = \text{Constant}$$

UCM

$$a_c = v$$

$$a_t = 0$$

### QUESTION 04

(Adv)

Match the matrix:


Column-I	Column-II
(a) UCM (r)(q) (p)	$\left  \frac{dv}{dt} \right  = 0$ <i>Magnitude of acc.</i>
(b) NUCM (r)(s) (q) $a_t \neq 0$	$\frac{d v }{dt} = 0$ <i>Magnitude of velocity = Const</i>
	(r) $\left  \frac{dv}{dt} \right  \neq 0$ $\rightarrow a_t$
	(s) $\frac{d v }{dt} \neq 0$



**QUESTION 05**


The motion of a particle will be circular if:

- A**  ~~$a_r = 0$  but  $a_t \neq 0$~~
- B**  ~~$a_r = 0$  and  $a_t = 0$~~
- C**  $a_r \neq 0$  but  $a_t = 0$  UCM
- D**  $a_r \neq 0$  and  $a_t \neq 0$  NUCM



**QUESTION 06**

The speed of a particle moving in a circle of radius  $r = 2$  m varies with time  $t$  as  $v = t^2$  where,  $t$  is in second and  $v$  in  $\text{ms}^{-1}$ . The net acceleration at  $t = 2$  s is [2012]



$$a_{\text{net}} = \sqrt{a_t^2 + a_c^2}$$

$$= \sqrt{8^2 + 4^2} = \sqrt{80}$$

Speed up.

$v = t^2 \Rightarrow t \rightarrow \text{vary } |v| \rightarrow \text{vary}$

NUCM.

$a_c = \frac{v^2}{r}$

$a_t = \frac{dv}{dt}$

$= \frac{d}{dt}(t^2) = 2t$

$a_c = \frac{4 \times 4}{2} = 8 \text{ m/s}^2$  at  $t = 2$

$a_t = 4 \text{ m/s}^2$



**QUESTION 07**

$K = \text{constant}$

$\alpha = -k\sqrt{\omega}$ , where  $\omega$  is the angular velocity of body. Find the time after which body will come to rest if at  $t = 0$ , angular velocity of body was  $\omega_0$ .

Sol  $\alpha = -k\sqrt{\omega}$   
 $\Downarrow$   
 vary as particle stops

$t=0 \quad \omega = \omega_0$   
 $t=t \quad \omega = 0$

ATDB.uno

$$\frac{d\omega}{dt} = -k\omega^{1/2}$$

$$\int_{\omega_0}^0 \frac{d\omega}{\omega^{1/2}} = \int_0^T -k dt \Rightarrow [2\sqrt{\omega}]_{\omega_0}^0 = -k[t]_0^T$$

$$T = \frac{2\sqrt{\omega_0}}{k}$$

**QUESTION 08**

(PYQ)

What is the value of linear velocity, if  $\vec{\omega} = 3\hat{i} - 4\hat{j} + \hat{k}$  and  $\vec{r} = 5\hat{i} - 6\hat{j} + 6\hat{k}$


Given: -  $\vec{v} = \vec{\omega} \times \vec{r}$

$$v = \begin{vmatrix} \hat{i} & -\hat{j} & \hat{k} \\ 3 & -4 & 1 \\ 5 & -6 & 6 \end{vmatrix}$$

$$= \underline{\hspace{2cm}}$$

### QUESTION 09

What is the average acceleration is going from A to B?




Total time.  
 $= \frac{R\sqrt{2}}{(\pi R/2)v}$

$$|\text{avg acc}| = \frac{|\vec{v}_f - \vec{v}_i|}{\Delta T}$$

$$= \frac{v\sqrt{2}}{(\pi R/2)/v}$$

Time taken =  $\frac{\pi R}{2v}$




$\vec{v}_f - \vec{v}_i =$

### QUESTION 10

$a = 15 \text{ m s}^{-2}$  represents the total acceleration of a particle, radius  $R = 2.5 \text{ m}$  at a given instant of time. The speed of the particle is UCM, NUCM

$a_c = a \cos 30$   
 $a_t = a \sin 30 \Rightarrow$  Speed up

$\frac{R}{v^2} = 15 \cos 30$   
 $v = \sqrt{R \times 15 \times \frac{1}{\cos 30}} = \sqrt{\frac{15}{2} \times 15 \times \frac{2}{\sqrt{3}}} = \dots$



ATDB.uno

**QUESTION 11**

CPYQ.

$\text{Power} = F \cdot v$

$= P = F v \cos \theta$



A particle of mass  $m$  is moving in a circular path of constant radius  $r$  such that its centripetal acceleration ( $a$ ) is varying with time  $t$  as  $a = k^2 r t^2$ , where  $k$  is a constant. The power delivered to the particle by the force acting on it is given as

- A** Zero
- B**  $mk^2 r^2 t^2$
- C**  $mk^2 r^2 t$
- D**  $mk^2 r t$

$\frac{v^2}{r} = a_c = k^2 r t^2$

NUCM


$v^2 = k^2 r^2 t^2$

$v = k r t$

$a_t = \frac{dv}{dt} = k r$

$F_c$  will not develop power.  
 $\theta = 90^\circ$

Power =  $F_t \cdot v \cos 0$   
 $= m a_t \times k r t$   
 $= m k r k r t$   
 $= m k^2 r^2 t$





**QUESTION 12**

(JEE)

$v = R\omega$

$A \rightarrow v_1 = R_1\omega$   
 $B \rightarrow v_2 = R_2\omega$

Two particles A and B are moving on two concentric circles of radii  $R_1$  and  $R_2$  with equal angular speed  $\omega$ . At  $t = 0$ , their positions and direction of motion are shown in the figure. The relative velocity  $\vec{v}_A - \vec{v}_B$  at  $t = \frac{\pi}{2\omega}$  is given by

**A**  $\omega(R_1 + R_2)\hat{i}$

**B**  $-\omega(R_1 + R_2)\hat{i}$

**C**  $\omega(R_1 - R_2)\hat{i}$

**D**  $\omega(R_2 - R_1)\hat{i}$   
 Ans

$\omega = \frac{2\pi}{T}$



$T = \frac{2\pi}{\omega}$

$\frac{T}{2} = \frac{\pi}{\omega}$

$t = \frac{1}{2} \left( \frac{T}{2} \right) = \frac{T}{4}$

ATDB.uno

$\vec{v}_A - \vec{v}_B$   
 $\Rightarrow -R_1\omega\hat{i} - (-R_2\omega)\hat{i}$   
 $= (-R_1\omega + R_2\omega)\hat{i}$   
 $= (R_2 - R_1)\omega\hat{i}$





# QUESTION 13

For a particle moving along circular path, the **radial** acceleration  $a_r$  is proportional to time  $t$ . If  $a_t$  is the tangential acceleration, then which of the following will be independent of time  $t$ ?

Centripetal Normal

**A**  $a_t$  ✗

**B**  $a_r \cdot a_t$  ✗

**C**  $a_r/a_t$

**D**  $a_r(a_t)^2$  ✓

$a_c \propto t$

$a_c = kt$  (NVCN)

$\frac{v^2}{R} = kt$

$v = \sqrt{Rkt}$

$a_t = \frac{dv}{dt} = \sqrt{Rt} \cdot \frac{1}{2\sqrt{t}}$

$a_r \cdot a_t \propto t \cdot \frac{1}{\sqrt{t}}$

$a_r \cdot a_t^2 \propto t \cdot \frac{1}{t}$

Independent of  $t$ .

ATDB.uno



**QUESTION 14**

(Level up)

$a_t = 0$   
UCM

A particle is describing uniform circular motion in the anti-clockwise sense such that its time period of revolution is  $T$ . At  $t = 0$  the particle is observed to be at  $A$ . If  $\theta_1$  be the angle between acceleration at  $t = \frac{T}{4}$  and average velocity in the time interval  $0$  to  $\frac{T}{4}$  and  $\theta_2$  be the angle between acceleration at  $t = \frac{T}{4}$  and the change in velocity in the time interval  $0$  to  $\frac{T}{4}$ , then


ATDB.uno

- A**  $\theta_1 = 135^\circ, \theta_2 = 45^\circ$
- B**  $\theta_1 = 135^\circ, \theta_2 = 135^\circ$
- C**  $\theta_1 = 45^\circ, \theta_2 = 135^\circ$
- D**  $\theta_1 = 45^\circ, \theta_2 = 45^\circ$






avg velocity =  $\frac{\text{Total disp}}{\text{Total time}}$



Change in velocity =  $\vec{v}_2 - \vec{v}_1$






## QUESTION 15

A rod  $AB$  is moving on a fixed circle of radius  $R$  with a constant velocity  $v$  as shown in figure.  $P$  is the point of intersection of rod and the circle. At an instant rod is at a distance  $x = \frac{3R}{5}$  from centre of circle. The velocity of rod is normal to its length and rod always remain parallel to the diameter  $CD$ . Find the speed of point  $P$  at this instant.

# ATDB.uno





# Circular Dynamics



## ❖ How to write force equation in Circular Motion.

# In CM

1. Draw FBD
2. Find Centre of Circle.
3. Resolve all forces along &  $\perp$  to Radius.


ATDB.uno



along radius  $F_{\text{net towards Centre}} = \frac{mv^2}{R} = mR\omega^2$

along Tangent  $F_{\text{net tangential}} = ma_t = mR\alpha$

Question 19



Pink thread:-


(Left Extreme)

Tension in thread =  $T_1$


(Right Extreme)

Tension in thread =  $T_2$

$$\frac{T_1}{T_2} = \frac{mg}{\cos \theta \times mg \cos \theta} = \frac{1}{\cos^2 \theta} = \sec^2 \theta$$



## ❖ Conical Pendulum



$$T \sin \theta = m R \omega^2$$

$$T \cos \theta = mg$$

$$\tan \theta = \frac{R \omega^2}{g}$$

$$\omega = \sqrt{\frac{g \tan \theta}{R}}$$


Time Period =  $\omega = \frac{2\pi}{T}$

$$\frac{2\pi}{T} = \sqrt{\frac{g \tan \theta}{R}}$$

$$T = 2\pi \sqrt{\frac{R}{g \tan \theta}}$$

$$R = l \sin \theta$$

## ❖ Bending of Cyclist



$$N \sin \theta = \frac{m v^2}{R}$$

$$N \cos \theta = mg$$

$$\tan \theta = \frac{v^2}{R g}$$

$$v = \sqrt{R g \tan \theta}$$

### # Banking of aeroplane / Train.


$$l \sin \theta = \frac{m v^2}{R}$$

$$l \cos \theta = mg$$

$$\tan \theta = \frac{v^2}{g r}$$




# ❖ Banking of Road



# + finding  $v_{max}$  for turn on banked road.

Car Tendency to Slip = outward.



ATDB.uno

$v \rightarrow v_{max}$   
 $f \rightarrow v_{max}$   
 $f = \mu_s N$

$$N \sin \theta + f \cos \theta = \frac{m v_{max}^2}{R}$$

$$N \cos \theta = mg + f \sin \theta$$

$$N (\sin \theta + \mu \cos \theta) = \frac{m v^2}{R}$$

$$N (\cos \theta - \mu \sin \theta) = mg$$

$\theta =$  Banking angle.



$$\frac{\sin\theta + \mu \cos\theta}{\cos\theta - \mu \sin\theta} = \frac{V_{\max}^2}{Rg}$$

Rough Road  
 but levelled Road  
 $\theta = 0$


$$V_{\max} \text{ for turn} = \sqrt{\frac{Rg (\sin\theta + \mu \cos\theta)}{(\cos\theta - \mu \sin\theta)}} = \sqrt{\frac{Rg (\tan\theta + \mu)}{1 - \mu \tan\theta}}$$

$$V = \sqrt{\mu Rg}$$

# What should be  $V_{\min}$  for banked Road

ATDB.uno

$$V_{\min} = \sqrt{\frac{Rg (\tan\theta - \mu)}{1 + \mu \tan\theta}}$$



$V_{\min} \Rightarrow$

$\mu \rightarrow -\mu$   
 Replace

# on banked Road  $\mu = 0$   $V = \sqrt{Rg \tan\theta}$

# ❖ Centrifugal Force



↓  
Pseudo force in CM.



ATDB.uno

Rest par maanege Earth ko

$$\frac{GMm}{R^2} = \frac{mv^2}{R}$$



$\frac{GMm}{R^2} = F_{\text{centripetal}}$  Inertial frame.

Isliye CM ho Raha hai.

$\frac{GMm}{R^2} = \frac{mv^2}{R}$



**QUESTION 16**

Three identical particles are joined together by a thread as shown in figure. All the three particles are moving on a smooth horizontal plane about point  $O$ . If the speed of the outermost particle is  $v_0$ , then the ratio of tensions in the three sections of the string is: (Assume that the string remains straight)

**A** 3 : 5 : 7

**B** 3 : 4 : 5

**C** 7 : 11 : 6

**D** 3 : 5 : 6


$\omega = \frac{\text{Angle Rotated}}{\text{Time}}$


ATDB.uno

for all three particles

$\omega = \frac{2\pi}{T} \Rightarrow \text{Same}$

$v = r\omega$   
 $v_3 = v_0 = 3l\omega$   
 $v_2 = 2l\omega$   
 $v_1 = l\omega$





$$F_{\text{net towards Centre}} = m\omega^2 R$$

$$= \frac{mv^2}{R}$$

ATDB.uno

$$T_1 = 6ml\omega^2$$

$$T_2 = 5ml\omega^2$$

$$T_3 = 3ml\omega^2$$

Ratio - 3:5:6.

~~$$T_1 - T_2 = ml\omega^2$$

$$T_2 - T_3 = m(2l)\omega^2$$

$$T_3 = m(3l)\omega^2$$~~

Add

$$T_1 = 6ml\omega^2$$

$$6ml\omega^2 - T_2 = ml\omega^2$$

$$5ml\omega^2 = T_2$$




# QUESTION 17

Goes particle has same speed at top  
 $WET \Rightarrow W_T = KE_f - KE_i$   
 $-mgH = \frac{1}{2}mv'^2 - \frac{1}{2}mv^2$        $v' = \text{Same in all cases.}$


A small block is shot into each of the four tracks as shown below. Each of the tracks rises to the same height. The speed with which the block enters the track is the same in all cases. At the highest point of the track, the normal reaction is maximum in

[2001] Adv.  
 ↳ Circular Motion




Least Radius  
 $R \downarrow$   
 $N \uparrow$

ATDB.uno



$$mg + N = \frac{mv'^2}{R}$$

$$N = \frac{mv'^2}{R} - mg$$



Contact force  $\rightarrow$  Normal  
 $\perp$  to surface  
 (Push) away.

Small cases  $v' \rightarrow$  Same

ATDB.uno  
 Radius = ?

Straight line  $\rightarrow$  part of circle whose  
 $R \rightarrow \infty$

Radius  $\downarrow$   $N \rightarrow$  Max.

More is the flatness  
 of surface  $R \uparrow$





## QUESTION 18

x 3 times

A bead of mass  $m$  stays at point  $P(a, b)$  on a wire bent in the shape of a parabola  $y = 4Cx^2$  and rotating with angular speed  $\omega$  (see figure). The value of  $\omega$  is (neglect friction)  
**[JEE (Main)-2020]**


**A**  $2\sqrt{2gC}$   
*Ans*

**B**  $2\sqrt{gC}$

**C**  $\sqrt{\frac{2g}{C}}$

**D**  $\sqrt{\frac{2gC}{ab}}$

ATDB.uno





ATDB.uno

$$N \sin \theta = m r \omega^2$$

$$N \cos \theta = mg$$

---


$$N \sin \theta = m a \omega^2$$

$$N \cos \theta = mg$$


---

$$\tan \theta = \frac{a \omega^2}{g}$$

$$\omega = \sqrt{\frac{g \tan \theta}{a}} = \sqrt{\frac{g \cdot 8c a}{a}} = 2\sqrt{2gc}$$

$$y = 4cx^2$$

$$\tan \theta = \frac{dy}{dx} = 8cx$$

at  $x = a$

$$\tan \theta = 8ca$$



## QUESTION 19

A modern grand-prix racing car of mass  $m$  is travelling on a flat track in a circular arc of radius  $R$  with a speed  $v$ . If the coefficient of static friction between the tyres and the track is  $\mu_s$ , then the magnitude of negative lift  $F_L$  acting downwards on the car is: (Assume forces on the four tyres are identical and  $g$  = acceleration due to gravity)

**[JEE (Main)-2021]**


**A**  $m \left( \frac{v^2}{\mu_s R} + g \right)$

**C**  $-m \left( g + \frac{v^2}{\mu_s R} \right)$

~~ATDB.uno~~


**B**  $m \left( \frac{v^2}{\mu_s R} - g \right)$

**D**  $m \left( g - \frac{v^2}{\mu_s R} \right)$





Tendency  
←



$\gamma \rightarrow$  Equilibrium

$$N = Mg + F$$


$$f = \frac{mv^2}{R}$$

$$MN = \frac{mv^2}{R}$$

ATDB.uno

$$M(Mg + F) = \frac{mv^2}{R}$$

$$F = \frac{mv^2}{MR} - Mg$$





**QUESTION 20**

HCV

In figure shows a rod of length 20 cm pivoted near an end and which is made to rotate in a horizontal plane with a constant angular speed. A ball of mass  $m$  is suspended by a string also of length 20 cm from the other end of the rod. If the angle  $\theta$  made by the string with the vertical is  $30^\circ$ , find the angular speed of the rotation. Take  $g = 10 \text{ m/s}^2$ .

$$T \cos 30 = mg$$



$$T \sin 30 = mR\omega^2$$


---


$$T \sin 30 = m(2 + 2 \sin 30)\omega^2$$

$$T \cos 30 = mg$$

Ratio  $\frac{1}{\sqrt{3}} = \frac{(3\frac{1}{2})\omega^2}{g}$   $\omega =$  \_\_\_\_\_





## QUESTION 21




A particle of mass  $m$  is fixed to one end of light spring having force constant  $k$  and unstretched length  $l_0$ . The other end is fixed. The system is given an angular speed  $\omega$  about the fixed end of the spring such that it rotates in a circle in gravity free space. Then the stretch in the spring is: **JEE Main 2020 – 8 Jan (Morning)**

**A**  $\frac{l_0 m \omega^2}{k - m \omega^2}$  *Ans*


**B**  $\frac{l_0 m \omega^2}{k + m \omega^2}$

**C**  $\frac{l_0 m \omega^2}{k - m \omega}$

**D**  $\frac{l_0 m \omega^2}{k + m \omega}$



ATDB.uno



ATDB.uno

$$F_s = -Kx$$



Restoring Nature.

Spring Elongate  
Compress

$F \rightarrow$  oppose  
 $F \rightarrow$  oppose.

$$F_s = m(l+x)\omega^2$$

$$Kx = m l \omega^2 + m x \omega^2$$

$$x(K - m\omega^2) = m l \omega^2$$

$$x = \frac{m l \omega^2}{K - m\omega^2}$$


**QUESTION 22**

(HCV)



A car goes on a horizontal circular road of radius  $R$ , the speed increasing at a constant rate  $\frac{dv}{dt}$ . The friction coefficient between the road and the tyre is  $\mu$ . Find the speed at which the car will skid.

Top View



$a_t = \frac{dv}{dt}$

When Car just slips

$f_{smax} = \mu N$

$f_{smax} = \mu mg$

$a_{net} = \sqrt{a_t^2 + a_c^2} = \sqrt{\left(\frac{dv}{dt}\right)^2 + \left(\frac{v^2}{R}\right)^2}$

$a_{net} = \sqrt{\left(\frac{dv}{dt}\right)^2 + \frac{v^4}{R^2}}$



$$F_{net} = m a_{net}$$

$$= m \sqrt{\left(\frac{dv}{dt}\right)^2 + \frac{v^4}{R^2}}$$

$$\text{frictional force} = F_{net} = \mu mg = m \sqrt{\left(\frac{dv}{dt}\right)^2 + \frac{v^4}{R^2}}$$

ATDB.uno

$$(\mu mg)^2 = \left(\frac{dv}{dt}\right)^2 + \frac{v^4}{R^2}$$

$$v = \sqrt[4]{R^2 \left[ (\mu mg)^2 - \left(\frac{dv}{dt}\right)^2 \right]}$$



## QUESTION 23

Two identical particles are attached at the ends of a light string which passes through a hole at the centre of a table. One of the particles is made to move in a circle on the table with angular velocity  $\omega_1$  and the other is made to move in a horizontal circle as a conical pendulum with angular velocity  $\omega_2$ . If  $l_1$  and  $l_2$  are the length of the string over and under the table, then in order that particle under the table neither moves down nor moves up the ratio  $l_1/l_2$  is:


**A**  $\frac{\omega_1}{\omega_2}$

**C**  $\frac{\omega_1^2}{\omega_2^2}$

**B**  $\frac{\omega_2}{\omega_1}$


**D**  $\frac{\omega_2^2}{\omega_1^2}$

ATDB.uno





$$T = ml_1 \omega_1^2$$




$$N = mg$$

ATDB.uno  
 $T \sin \theta = mr \omega_2^2$

$$ml_1 \omega_1^2 \sin \theta = m l_2 \sin \theta \omega_2^2$$

$$\frac{l_2}{l_1} = \frac{\omega_2^2}{\omega_1^2}$$



$$T \cos \theta = mg$$



## QUESTION 24

A smooth wire of length  $2\pi r$  is bent into a circle and kept in a vertical plane. A bead can slide smoothly on the wire. When the circle is rotating with angular speed  $\omega$  about the vertical diameter  $AB$ , as shown in figure, the bead is at rest with respect to the circular ring at position  $P$  as shown. Then the value of  $\omega^2$  is equal to

**[JEE (Main)-2019]**

**A**  $\frac{(g\sqrt{3})}{r}$

**C**  $\frac{2g}{(r\sqrt{3})}$

**B**  $\frac{2g}{r}$

**D**  $\frac{\sqrt{3g}}{2r}$





## QUESTION 25

A uniform ring, having radius  $a$  and mass  $m$  is to be rotated in the horizontal plane about its own axis with constant angular velocity  $\omega$ . What would be the tension in the ring and nature of force?

- A**  $\frac{MR\omega^2}{2\pi}$  tensile
- B**  $MR\omega^2$  tensile
- C**  $\frac{MR\omega^2}{2}$  compressive
- D**  $MR\omega^2$  compressive

ATDB.uno




## QUESTION 26


(HCV)

A table with smooth horizontal surface is placed in a cabin which moves in a circle of a large radius  $R$ . A smooth pulley of small radius is fastened to the table. Two masses  $m$  and  $2m$  placed on the table are connected through a string going over the pulley. Initially the masses are held by a person with the strings along the outward radius and then the system is released from rest (with respect to the cabin). Find the magnitude of the initial acceleration of the masses as seen from the cabin and the tension in the string.

ATDB.uno

Top View.





ATDB.uno

$$2mR\omega^2 - T = 2ma$$

$$T - mR\omega^2 = ma$$

$$mR\omega^2 = 3ma$$

$$a = \frac{R\omega^2}{3}$$

**QUESTION 27**

HCV





A table with smooth horizontal surface is fixed in a cabin that rotates with a uniform angular velocity  $\omega$  in a circular path of radius  $R$ . A smooth groove  $AB$  of length  $L$  ( $\ll R$ ) is made on the surface of the table. The groove makes an angle  $\theta$  with the radius  $OA$  of the circle in which the cabin rotates. A small particle is kept at the point  $A$  in the groove and is released to move along  $AB$ . Find the time taken by the particle to reach the point  $B$

gadha.

Force along groove =  $mR\omega^2 \cos\theta = m/a$  ATDB.uno

$a = R\omega^2 \cos\theta$



$MR\omega^2 = f_{centrifugal}$



$$s = ut + \frac{1}{2} at^2$$

$$a = R\omega^2 \cos \theta$$

Constant

ATDB.uno

$$h = \frac{1}{2} (R\omega^2 \cos \theta) t^2$$

$$\sqrt{\frac{2h}{R\omega^2 \cos \theta}} = t$$



## QUESTION 28

A ball of mass ( $m$ ) 0.5 kg is attached to the end of a string having length ( $L$ ) 0.5 m. The ball is rotated on a horizontal circular path about vertical axis. The maximum tension that the string can bear is 324 N. The maximum possible value of angular velocity of ball (in radian/s)

**[JEE 2011]**


$$T_{\max} = 324 \text{ N.}$$

$$T \sin \theta = m r \omega^2$$

$$T \sin \theta = m L \sin \theta \omega^2$$

$$T_{\max} = m L \omega_{\max}^2$$

$$\omega = \sqrt{\frac{324}{0.5 \times 0.5}} = \frac{18}{0.5} = 36$$











# QUESTION 29

Two identical discs of same radius  $R$  are rotating about their axes in opposite directions with the same constant angular speed  $\omega$ . The disc are in the same horizontal plane. At time  $t = 0$ , the points  $P$  and  $Q$  are facing each other as shown in the figure. The relative speed between the two points  $P$  and  $Q$  is  $v_r$ . as function of times best represented by

**[IIT-JEE-2012]**





$t = 0$	0
$t = T/4$	$2v$
$t = T/2$	0
$t = 3T/4$	$2v$
$t = T$	0

ATDB.uno

$$V_{RQ} = V_P - V_Q$$

$$= -v - (+v)$$

$$= -2v$$

Relative speed =  $2v$ .

$t=0$

$$V_r = 0$$





## QUESTION 30

A wire, which passes through the hole is a small bead, is bent in the form of quarter of a circle. The wire is fixed vertically on ground as shown in the figure. The bead is released from near the top of the wire and it slides along the wire without friction. As the bead moves from A to B, the force it applies on the wire is **[JEE Adv 2014]**

- A** always radially outwards
- B** always radially inwards
- C** radially outwards initially and radially inwards later
- D** radially inwards initially and radially outwards later.


ATDB.uno





ATDB.uno

$N \rightarrow$  Force on bead by wire.



$$mgR(1 - \cos \theta) = \frac{1}{2}mv^2 - 0$$

$$v =$$

$$mg \sin \theta = F_t = ma_t$$

ATDB.uno

Q at what  $\theta$  block leaves Contact.

$$mg \cos \theta - N = \frac{mv^2}{R}$$

$$N=0$$

$$\theta = \cos^{-1} \left( \frac{2}{3} \right)$$






## QUESTION 31

2022  
(Advanced PYQ)

List-I describes four systems, each with two particles  $A$  and  $B$  in relative motion as shown in figure. List-II gives possible magnitudes of their relative velocities (in  $\text{ms}^{-1}$ ) at time  $t = \frac{\pi}{3}$  s.

$\vec{V}_A - \vec{V}_B$

List-I		List-II	
I.	<p><math>A</math> and <math>B</math> are moving on a horizontal circle of radius 1 m with uniform angular speed <math>\omega = 1 \text{ rad s}^{-1}</math>. The initial angular positions of <math>A</math> and <math>B</math> at time <math>t = 0</math> are <math>\theta = 0</math> and <math>\theta = \frac{\pi}{2}</math> respectively.</p>  <p><math>v = r\omega = 1 \times 1 = 1 \text{ m/s}</math></p>	P.	$\frac{\sqrt{3} + 1}{2}$



ATDB.uno


$$V_{rel} = \sqrt{1^2 + 1^2 - 2 \cdot 1 \cdot 1 \cdot \cos 90} = \sqrt{2}$$

$t = \pi/3$   
 angle rotated,  $\theta = \omega t + \frac{1}{2} \alpha t^2$   
 $= 1 \times \frac{\pi}{3}$   
 $= 60^\circ$

## List-I

## List-II


- II. Projectiles  $A$  and  $B$  are fired (in the same vertical plane) at  $t = 0$  and  $t = 0.1$  s respectively, with the same speed  $v = \frac{5\pi}{\sqrt{2}} \text{ ms}^{-1}$  and at  $45^\circ$  from the horizontal plane. The initial separation between  $A$  and  $B$  is large enough so that they do not collide, ( $g = 10 \text{ ms}^{-2}$ ).



$$V_{rel} \text{ at } t = \frac{\pi}{3}$$

ATDB.uno


- III. Two harmonic oscillators  $A$  and  $B$  moving in the  $x$  direction according to  $x_A = x_0 \sin \frac{t}{t_0}$  and  $x_B = x_0 \sin \left( \frac{t}{t_0} + \frac{\pi}{2} \right)$  respectively, starting from  $t = 0$ . Take  $x_0 = 1$  m,  $t_0 = 1$  s.



R.

$$\sqrt{10}$$





$$\vec{V}_A = \frac{5\pi}{2} \hat{i} + \left(\frac{5\pi}{2} - 1\right) \hat{j}$$

$$\vec{V}_B = -\frac{5\pi}{2} \hat{i} + \frac{5\pi}{2} \hat{j}$$

$$\vec{V}_A - \vec{V}_B = 5\pi \hat{i} - \hat{j}$$

$$|\vec{V}_A - \vec{V}_B| = \sqrt{25\pi^2 + 1}$$

$$V_y = u_y + a_y t$$

$$= \frac{5\pi}{2} - 10 \times 0.1$$

$$= \frac{5\pi}{2} - 1$$

$a_{rel} = 0$

ATDB.uno

vel. rel. appas mein Same after any time.

$t = \pi/3$  nikalo  
ya  $t = 0.1$  sec  
par nikalo  
baad ek  
hi hai.




### List-I

### List-II

IV.

Particle A is rotating in a horizontal circular path of radius 1 m on the xy plane, with constant angular speed  $\omega = 1 \text{ rads}^{-1}$ . Particle B is moving up at a constant speed  $3 \text{ ms}^{-1}$  in the vertical direction as shown in the figure. (Ignore gravity.)



$v = r\omega$   
 $v = 1$

$v_B = \hat{z}$   
 $v_A = \hat{i} \text{ or } \hat{j} \text{ or } \hat{i} + \hat{j}$   
 $v_{rel} = \sqrt{3^2 + 1^2} = \sqrt{10}$

S.

$\sqrt{2}$

T.

$\sqrt{25\pi^2 + 1}$

ATDB.uno

Which one of the following options is correct?

(A) I → R, II → T, III → P, IV → S  
(B) I → S, II → P, III → Q, IV → R  
(C) I → S, II → T, III → P, IV → R  
(D) I → T, II → P, III → R, IV → S

~~(A)~~  
~~(B)~~  
~~(D)~~  
✓  
✓  
✓  
Ans

I - (S)



## QUESTION 32

(HW)

A long horizontal rod has a bead which can slide along its length and is initially placed at a distance  $L$  from one end  $A$  of the rod. The rod is set in angular motion about  $A$  with a constant angular acceleration  $\alpha$ . If the coefficient of friction between the rod and bead is  $\mu$ , and gravity is neglected, then the time after which the bead starts slipping is **(2000, 2M)**

**A**  $\sqrt{\frac{\mu}{\alpha}}$

ATDB.uno

**B**  $\frac{\mu}{\sqrt{\alpha}}$



**C**  $\frac{1}{\sqrt{\mu\alpha}}$

**D** infinitesimal


## ❖ Radius of curvature



Circle is a Geometrical shape which has same Radius at all points.



ATDB.uno



for other Curves  $y = f(x)$

Radius will be different at different Points.

ROI BHI mathematical eqn dekh


#



$y = f(x)$

$$R = \frac{\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{3/2}}{\left| \frac{d^2y}{dx^2} \right|}$$

slope



ATDB.uno

$$y = x \tan \theta - \frac{gx^2}{2u^2 \cos^2 \theta}$$

$$R_{\text{at top}} = \frac{\left[ 1 + (0)^2 \right]^{3/2}}{\left| \frac{-g}{u^2 \cos^2 \theta} \right|} = \frac{u^2 \cos^2 \theta}{g}$$

slope  $\Rightarrow \frac{dy}{dx} = \tan \theta - \frac{gx}{u^2 \cos^2 \theta}$  Put  $x = \frac{R}{2}$

$$\frac{d^2y}{dx^2} = 0 - \frac{g}{u^2 \cos^2 \theta}$$

#! a straight line  $\kappa \rightarrow \infty$

$$y = mx + c$$


$$\frac{dy}{dx} = m$$

$$\frac{d^2y}{dx^2} = 0$$

$$R = \frac{[1 + m^2]^{3/2}}{0}$$



radius infinity



ATDB.uno


$$g \Rightarrow a_{centripetal} = \frac{v^2}{r}$$

$$g = \frac{u^2 \cos^2 \theta}{R}$$

$$R = \frac{u^2 \cos^2 \theta}{g}$$



#



Find R of Curvature when particle makes angle  $\theta/2$  with horizontal.

$$\frac{dy}{dx} = \tan(\theta/2)$$

$$\left| \frac{d^2y}{dx^2} \right| = \frac{g}{u^2 \cos^2 \theta}$$

$$R = \frac{\left[ 1 + \tan^2(\theta/2) \right]^{3/2}}{\frac{g}{u^2 \cos^2 \theta}}$$

**QUESTION 33**

(Irodov)

A particle moves at uniform speed on a parabolic trajectory  $y = ax^2$  at uniform speed  $v$ . Find the acceleration of particle when it passes point  $x = 0$  and point  $(1, a)$ .

ATDB.uno



Thank  
YOU

ATDB.uno



Keep Hustling!

