

ARJUNA

JEE AIR 2024

ATDB.uno
Physical Chemistry

Ionic Equilibrium



Lecture No. - 08

**By- Dr. Gopal Chaturvedi Sir
(GC Sir)**

**GURUKUL**

Topics to be Covered

Topic

Titration

Topic

Indicator Theory

TopicSolubility (S) **ATDB.uno****Topic**Solubility product (K_{SP})



GURUKUL

Recap of Previous Lecture


Topic

Buffer Action

ATDB.uno

Dilution of Buffer solution :-

Addition of $H_2O(l)$




Before dilution ;

$$K_a = \frac{(C_1 + x)x}{(C_2 - x)} ; x = [H^+]$$

After dilution ;

$$C_1 \& C_2 \quad K_a = \frac{(C'_1 + x')x'}{(C'_2 - x')} ; x' = [H^+]_{new}$$

After dilution.



(i). Addition of $H_2O(l)$ in small amount ; no appreciable change in pH


(ii). Addition of $H_2O(l)$ in large amount ; pH of buffer solution $\rightarrow 7$ at $25^\circ C$

**GURUKUL**

In general indicators are either weak organic acids or weak organic bases having different coloured characteristics in ionised and unionised form.

General acidic indicator ; HIn

ATDB.uno



$$K_{\text{HIn}} \text{ or } K_{\text{In}} = \frac{[\text{H}^+]_{\text{sol}} [\text{In}^-]}{[\text{HIn}]}$$

ATDB.uno

$$[\text{H}^+]_{\text{sol}} = K_{\text{In}} \left\{ \frac{[\text{HIn}]}{[\text{In}^-]} \right\}$$



GURUKUL

$$\boxed{\text{pH} = \text{pK}_{\text{In}} + \log \frac{[\text{In}^-]}{[\text{HIn}]}} \quad \checkmark$$

$$\% \text{ ionised form of indicator} = \frac{[\text{In}^-]}{[\text{HIn}] + [\text{In}^-]} \times 100 = \frac{[\text{In}^-] / [\text{HIn}]}{1 + \frac{[\text{In}^-]}{[\text{HIn}]}} \times 100$$

$$\% \text{ unionised form of indicator} = \frac{[\text{HIn}]}{[\text{HIn}] + [\text{In}^-]} \times 100$$

ATDB.uno




Titration



GURUKUL

This is an experimental technique by which we can determine amount of unknown concⁿ solution with the help of reaction between different nature of substance.

ATDB.uno



Standard Solution (Titrant)
[known concⁿ solution]

M_2 e.g. Acid $HCl(aq)$


ATDB.uno

Analyte [Unknown concⁿ solution]

V_1 ml (given)

M_1 (Unknown)

e.g. $NaOH(aq)$



GURUKUL

Given.

 M_2 $V_1 \text{ ml}$ $V_2 \text{ ml}$ M_1

(Used)

(Unknown)

When $M_1 V_1 = M_2 V_2$ Equivalence point or neutralisation point.

 NaCl(aq) Sol^n

$$M_1 = \frac{M_2 V_2}{V_1}$$



NaOH base

M_2, V_2

$\text{CH}_3\text{COOH}(\text{acid})$

M_1, V_1

ATDB.uno

At equivalent point in acid-base neutralisation
No. of equivalent of Acid = No. of equivalent of base


$$1 \times \boxed{M_1 V_1} = \boxed{M_2 V_2} \times 1$$

Indicator change its colour in certain pH range, this pH range is called colour transition range.



$$\frac{[\text{In}^-]}{[\text{HIn}]} \approx \frac{1}{10} \text{ to } \frac{10}{1}$$


$$\boxed{\text{pH} = \text{pKIn} \pm 1}$$



^{pH}
ATDB.uno
Lower

^{pH}
Higher

General basic indicator : InOH




ATDB.uno



GURUKUL

Methyl Orange : Basic Indicator



⇒ If pH of solution ≥ 4.4 ; then solution has yellow colour.

⇒ ✓ If pH of solution ≤ 3.1 ; then solution has (red) colour


⇒ In between 3.1 to 4.4 ; indicator change its colour.

ATDB.uno



GURUKUL

Phenolphthalein :- Acidic indicator



ATDB.uno

⇒ If pH of solution ≥ 10 ; then solution has pink colour.

⇒ If pH of solution ≤ 8.2 ; then solution is colourless.

⇒ In between 8.2 to 10 ; indicator change its colour.

CTR
↓
(8.2 → 10)
✓



Q.

Titration of SA with SB

Q. If $HCl(aq)$ is titrated against $0.1M NaOH(aq)$ solution.
(50ml)

Used Volume of $NaOH$ (1) 0 ml

ATDB.uno

(2) 49 ml

(3) 50 ml


Calculate the p^H of solⁿ. (4) 51 ml

(1) Resulting solⁿ.

HCl(aq)
50ml
0.1M

$$P^H = -\log 10^{-1} = 1$$

ATDB.uno





Resulting Solution HCl(aq) & NaCl(aq)

$$(\text{HCl(aq)}) = \frac{(5 - 4.9)}{99} = \frac{0.1}{99}$$

$$P^H = -\log\left(\frac{0.1}{99}\right) \approx 3$$



GURUKUL



$\longrightarrow \text{p}^{\text{H}} = \frac{1}{2} \text{p}^{\text{Kw}} = 7$ at 25°C

Neutralisation point.



After reaction mmole — (5.1 - 5) = 0.1



$[\text{NaOH(aq)}] = \frac{0.1}{101} \approx 10^{-3}$
 $\text{p}^{\text{OH}} \approx 3, \text{p}^{\text{H}} = 11$ at 25°C

Titration of WA against SB



If $\text{CH}_3\text{COOH(aq)}$ is titrated against 0.1M NaOH(aq) solution
(50ml)
(0.1M)

Used volume of 0.1M NaOH(aq) are

(1) 0 ml

(2) 25 ml

(3) 49 ml

(4) 50 ml

(5) 51 ml

$$K_a = 1.8 \times 10^{-5}$$

(CH_3COOH)

ATDB.uno


Calculate p^{H} of resulting solution



GURUKUL

Sol.
1.

Resulting Solution




W. Acid Solⁿ

$$[H^+] = \sqrt{K_a \times C}$$


$$p^H = \frac{1}{2}(p^{K_a} - \log c)$$
$$= \frac{1}{2}(4.74 + 1) = 2.87$$

ATDB.uno

$$p^{K_a} = -\log(1.8 \times 10^{-5})$$
$$= 4.74$$



After reactⁿ mmoles 2.5



$p^H = p^{K_a}$
 $p^H = 4.74$
Max. Buffer Capacity
⑤ = ④



mmoles 5 4.9


mmoles 0.1 — 4.9 4.9

Resulting solution $\text{CH}_3\text{COOH}(\text{aq})$, $\text{CH}_3\text{COONa}(\text{aq})$



ATDB.uno

$$\begin{aligned} \text{pH} &= 4.74 + \log\left(\frac{4.9}{0.1}\right) = 4.74 + 2 \log 7 \\ &= 6.43 \end{aligned}$$

log 7
 ↓
 0.8451



GURUKUL




Basic ✓ $[\text{CH}_3\text{COONa}] = \frac{5}{100} = 0.05$

$p^H = p^{Kw} - p^{OH}$, $p^H = \frac{1}{2} p^{Kw} + \frac{1}{2} (p^{Ka} + \log C)$

$= 7 + \frac{1}{2} [4.74 + \log(0.05)]$

≈ 8.71




$[\text{NaOH}] = \frac{0.1}{101} = c_1$

ATDB.uno

Hydrolysis of CH_3COO^- in pr. of

SB




$[\text{OH}^-]_{\text{T}} = [\text{OH}^-]_{\text{from NaOH}} + x \approx c_1$

$[\text{OH}^-] = 10^{-3}$
 $\text{pH} = 11$



GURUKUL



Question



GURUKUL

- #Q. (a) Determine the pH of a 0.2 M solution of pyridine C_5H_5N . $K_b = 1.5 \times 10^{-9}$
- (b) Predict the effect of addition of pyridinium ion $C_5H_5NH^+$ on the position of the equilibrium. Will the pH be raised or lowered?
- (c) Calculate the pH of 1.0 L of 0.10 M pyridine solution to which 0.3 mol of pyridinium chlorides $C_5H_5NH^+ Cl$, has been added, assuming no change in volume.

HW

ATDB.uno





GURUKUL

Ans. (a) $\text{pH} = 9.239$ (b) lowered (c) $\text{pH} = 4.699$

ATDB.uno

Question



GURUKUL

#Q. Calculate the pH of a solution which result from the mixing of 50.0 ml of 0.3 M HCl with 50.0 ml of 0.4 M NH_3 . [$K_b (\text{NH}_3) = 1.8 \times 10^{-5}$]

HW

ATDB.uno

Ans. 8.7782



ATDB.uno

Question



GURUKUL

#Q. Calculate the pH of a solution made mixing 50.0 ml of 0.2 M NH_4Cl & 75.0 ml of 0.1 M NaOH . [$K_b(\text{NH}_3) = 1.8 \times 10^{-5}$]

HW

ATDB.uno

Ans. 9.7324



ATDB.uno

Question



GURUKUL

#Q. A buffer solution was prepared by dissolving 0.02 mol propionic acid & 0.015 mol sodium propionate enough water to make 1.00 L of solution (K_a for propionic acid is 1.34×10^{-5})

- (a) What is the pH of the buffer?
- (b) What would be the pH if 1.0×10^{-5} mol HCl were added per 10 ml of the buffer?
- (c) What would be the pH if 1.0×10^{-5} mol NaOH were added per 10 ml of the buffer.

ATDB.uno

HW



GURUKUL

Ans. (a) 4.7525 (b) 4.697, (c) 4.798

ATDB.uno

Question

#Q. A solution was made up to be 0.01 M in chloroacetic acid, ClCH_2COOH and also 0.002 M in sodium chloroacetate $\text{ClCH}_2\text{COONa}$. What is $[\text{H}^+]$ in the solution? $K_a = 1.5 \times 10^{-3}$.

HW

ATDB.uno




Ans. $[H^+] = 2.5 \times 10^{-3}$



ATDB.uno

Question



GURUKUL

#Q. For the indicator thymol blue, pH is 2.0 when half the indicator is in unionized form. Find the % of indicator in unionized form in the solution with $[H^+] = 4 \times 10^{-3} \text{ M}$.

HW

ATDB.uno

Ans. $[HI_n] = 28.57\%$



ATDB.uno

Question

#Q. Bromophenol blue is an indicator with a K_a value of 6×10^{-5} . What % of this indicator is in its basic form at a pH of 5?

HW

ATDB.uno



Ans. 85.71%



ATDB.uno

Question



GURUKUL

#Q. An acid base indicator has a K_a of 3×10^{-5} . The acid form of the indicator is red & the basic form is blue. By how much must the pH change in order to change the indicator form 75% red to 75% blue?

HW

ATDB.uno

Ans. $\Delta\text{pH} = 0.954$



ATDB.uno

Question



GURUKUL

#Q. Calculate the hydronium ion concentration and pH at the equivalence point in the reaction of 22.0 mL of 0.10 M acetic acid, CH_3COOH , with 22.0 mL of 0.10 M NaOH .

HW

ATDB.uno

Ans. 8.71



ATDB.uno

Question

#Q. Calculate the hydronium ion concentration and the pH at the equivalence point in a titration of 50.0 mL of 0.40 M NH_3 with 0.40 M HCl.

HW


ATDB.uno



Ans. 4.98



ATDB.uno



GURUKUL



Thank You ATDB.uno

