

PRAAYAS

JEE 2026

ATDB.uno

PHYSICAL CHEMISTRY

REDOX REACTION

Lecture – 02

FAISAL RAZAQ





Topics to be covered

- A** Concept of Oxidation and Reduction
- B** Oxidation State

ATDB.uno






TELEGRAM GROUP BY FAISAL SIR



ATDB.uno





Khoobsurat Baat 😊

Yield = $\frac{\text{Jitna bana}}{\text{Jitna banana chahiye tha}} \times 100\%$

reaction likho lekin balance karna
bhul gye toh bech bilkul garg 🤪

A compound H_2X with molar weight of 80 g is dissolved in a solvent having density of 0.4 $g\ mL^{-1}$. Assuming no change in volume upon dissolution, the molality of a 3.2 molar solution is

Sol. $M = 3.2 = \frac{n_b}{V_{\text{solution}}}$

$W_{\text{Solvent}} = 0.4 \times 1000 = 400g = W_A$

3.2 moles of H_2X in 1000 ml of Solution

Molality = $\frac{n_b}{W_A} \times 1000 = 3$

[JEE Adv. -2014]

104. Number of grams of bromine that will completely react with 5.0 g of pent-1-ene is $\times 10^{-2}$ g. (Atomic mass of Br = 80 g/mol) (Nearest integer)

[JEE Mains 25 June, 2022, Shift-I]

First write the reaction

CCCC=C + Br2 -> CCCC(Br)C(Br)C ($C_5H_{10}Br_2$)

Moles of $C_5H_{10} = 1143 \times 10^{-2}$ g

dimag ki jali batti 🤪

adv mei aaya hai, ignore karne
ki galti mat karna 🤪

143. The mole fraction of urea in aqueous urea solution containing 900 g of water is 0.05. If the density of the solution is $1.2\ g\ cm^{-3}$, the molarity of urea solution is ____.

(Given data: Molar masses of urea and water are $60\ g\ mol^{-1}$ and $18\ g\ mol^{-1}$, respectively)

[JEE Adv. -2019]

Sol. $X_{\text{urea}} = 0.05 = \frac{n}{n+50}$, $19n = 50$, $n = 2.6315$

$V_{\text{sol}} = \frac{(2.6315 \times 60 + 900)}{1.2} = 881.5789\ ml$

Molarity = $\frac{2.6315 \times 1000}{881.5789} = 2.9849$

NO₂

0.5 of nitrogen atom = +3

0.5 of both oxygen atoms = -2

K₂Cr₂O₇

0.5 of chromium atom = +6

0.5 of all oxygen atoms = -2

CCl₄

0.5 of carbon atom = +4

0.5 of all chlorine atoms = -1

[O-]N(=O)[O-]

[K+].[O-]C(=O)C(=O)[O-].[K+]

ClC(Cl)(Cl)Cl


Common Mistake 🤪

Avg. at. wt. = $(54 + 56 + 57)/3$

This is possible only when all the isotopes are in same percentage.

n-factor Calculation for acids



$$\text{Eq. wt} = \frac{\text{Molar wt}}{n}$$



$E_2 = \text{Humme nahi pata!}$

if the equivalent weight of $K_x H_y (C_2O_4)_z$ is E_1 and E_2 then find out the value of E_1 and E_2 in terms of M .

#



$E_2 = \text{Hume wahi pata!}$

ATDB.uno

Question

(F)

S1: The n-factor of acid is always equal to its basicity
(True/False)

(F)

S2: The n-factor of HCl is always 1
(True/False)

A) T, T

B) T, F

C) F, F

D) F, T

Sometimes HCl behaves like a
reducing agent.

n-Factor Calculation for Base




replaceable $\bar{O}H$ (hydroxyl ions)

or

acceptable H^+ ions

	Acidity	n-factor
$NaOH \rightarrow$	1	1
$Ca(OH)_2 \rightarrow$	2	1, 2
$Al(OH)_3 \rightarrow$	3	1, 2, 3



ATDB.uno

N-factor Calculation for Salts



- i)
 - ii)
 - iii)
 - ...
- } Oxidation state


ATDB.uno



Oxidation State

"The oxidation state of an atom can be defined as the hypothetical charge that would be held by that atom if all of its bonds to other atoms were completely ionic in nature."


ATDB.uno



O.S of O atom = -2



O.S of both H-atoms = +1

ATDB.uno




O.S of both O-atoms = -1


O.S of both H-atoms = +1








$$\left. \begin{array}{l} S = +6 \\ H = +1 \\ O = -2 \end{array} \right\}$$




$$Mn = +7$$



$$Cl = +7$$




Carbon Suboxide (C_3O_2)



among 3-carbons $\begin{cases} 2C = +2 \\ 1C = 0 \end{cases}$


$$\text{Avg O.S. of C} = \frac{2+0+2}{3} = +\frac{4}{3}$$

Cr_2O_5 (Butterfly str)




O.S of Cr = +6
Among 5 O atoms $\begin{cases} 4O = -1 \\ 1O = -2 \end{cases}$

Caro's Acid (H_2SO_5)



O.S of S atom = +6
 Among 5-O-atoms $\left\{ \begin{array}{l} 3O = -2 \\ 2O = -1 \end{array} \right.$


Marshall's Acid ($H_2S_2O_8$)



Among 2 S atoms both are +6
 Avg = $\frac{+6 + 6}{2} = +6$
 Among 8 O-atoms $\left\{ \begin{array}{l} 6O = -2 \\ 2O = -1 \end{array} \right.$


ATDB.uno

Tribromooxide (Br_3O_8)



$$\text{Avg}_O = \frac{+6 + 4 + 6}{3} = +\frac{16}{3}$$

Tetrathionate ion ($\text{S}_4\text{O}_6^{2-}$)




$$\text{Avg}_O = \frac{5 + 0 + 0 + 5}{4} = \frac{5}{2} = 2.5$$

ATDB.uno

JEE Adv. 2011 Paper 1



The difference in the oxidation states of two types of sulphur atoms in $\text{Na}_2\text{S}_4\text{O}_6$ is



Rules governing oxidation state



i) Fluorine Atom

* Fluorine is most electronegative element. It always has oxidation number equal to -1 in all its compounds.

* In case of F_2 it is zero.





ii) Oxygen Atom



a) In general as well as in its oxides = -2

b) In peroxide (H_2O_2 , Na_2O_2 etc) = -1

c) In superoxide (KO_2 etc) = $-\frac{1}{2}$

d) In ozonide (KO_3 etc) = $-\frac{1}{3}$

ATDB.uno

e) In O_2F_2 and OF_2 = +1, +2 respectively

f) In oxygen molecule (O_2) = 0

III) Hydrogen Atom



* In general hydrogen atom has oxidation number equal to +1.

* In metallic hydrides (NaH, KH etc) it is -1.

* In H_2 the O.S of H = 0


ATDB.uno

iv) Halogen Atom




* In general halogen (Cl, Br, I) has oxidation number = -1

* When halogen is attached with more electronegative element then it will show positive oxidation state.



ATDB.uno



V) Metals



* Alkali metals (Li, Na, K, Rb - - -) = +1

* Alkaline earth metals (Be, Mg, Ca, Sr - - -) = +2

* Aluminium always has +3 oxidation number.

ATDB.uno

vi) Oxidation number of an element in free state or in allotropic forms is always zero. $S_8, P_4, Fe_{(s)}, Na_{(s)}$

vii) Sum of charges of elements in a molecule is zero.

ATDB.uno

viii) Sum of charges of all elements in an ion is equal to the charge on that ion.

LIST OF IONS

H.Wo

Cl⁻	chloride	C₂O₄²⁻	oxalate
Br⁻	bromide	NO₃⁻	nitrate
F⁻	fluoride	N³⁻	nitride
I⁻	iodide	NO₂⁻	nitrite
CO₃²⁻	carbonate	ClO₄⁻	perchlorate
CN⁻	cyanide	ClO₃⁻	chlorate
NC⁻	isocyanide	ClO₂⁻	chlorite
SO₄²⁻	sulphate	ClO⁻	hypochlorite
SO₃²⁻	sulphite	CrO₄²⁻	chromate
S₂O₃²⁻	thiosulphate	Cr₂O₇²⁻	dichromate
S²⁻	sulphide	MnO₄⁻	permanganate
P³⁻	phosphide	PO₄³⁻	phosphate

ATDB.uno



$$2(+1) + x + 4(-2) = 0$$

$$x = +6$$



$$+1 + x + 4(-2) = 0$$

$$x = +7$$

ATDB.uno



$$3x + 8(-2) = 0$$

$$x = +\frac{16}{3}$$

Question



If three electrons are lost by a metal ion M^{3+} , its final oxidation number should be :

- A 0
- B +6
- C +2
- D +4

ATDB.uno

Question



Oxidation number of Sulphur in Na_2SO_4 is

A -2

B +6

C +2

D -6

ATDB.uno

JEE Main 2 Sep 2022 Shift-2



The oxidation states of transition metal atoms in $K_2Cr_2O_7$, $KMnO_4$ and K_2FeO_4 , respectively, are x , y and z . The sum of x , y and z is _____.

ATDB.uno

Question



Oxidation number of Sulphur in S_2Cl_2 is :

A +1

B 0

C -1

D +6

ATDB.uno

JEE Main 2019



N_2O_3 , NO , NO_2 , N_2O arrange the increasing order of O.S. of nitrogen –

- A N_2O , NO , N_2O_3 , NO_2
- B N_2O , NO_2 , N_2O_3 , NO
- C NO_2 , N_2O_3 , N_2O , NO
- D N_2O_3 , NO , N_2O , NO_2

ATDB.uno

Question



In which of the following compound oxidation number of Cl is +3?

- A ICl
- B ClO_3
- C ClF_3
- D HClO_4

ATDB.uno

Question



Which of the following is the correct oxidation number of phosphorus in $\text{Mg}_2\text{P}_2\text{O}_7$

A -3

B +2

C +5

D +3

ATDB.uno

Question



Oxidation numbers of the two nitrogen atoms present in ammonium nitrate are respectively ?

- A +3 and +3
- B 0 and 0
- C -3 and +5
- D -1 and -1

ATDB.uno

Question



The oxidation number of cobalt in $[\text{Co}(\text{CN})_6]^{3-}$ is –

A +3

B -3

C +6

D -6

ATDB.uno

Question



Oxidation number of Sulphur in $S_2O_2^{2-}$ is :

A -2

B +1

C +6

D 0

ATDB.uno

Question



In which of the following compound oxidation number of iron is not +3

- A Fe_3O_4
- B Fe_2O_3
- C FeCl_3
- D FePO_4

ATDB.uno

Question



In which of the following compound oxidation number of iron is not +3

- A Fe_3O_4
- B Fe_2O_3
- C FeCl_3
- D FePO_4

ATDB.uno

Question



Oxidation number of Sulphur in H_2SO_5 is

A +2

B +4

C +8

D +6

ATDB.uno

Question



Oxidation number of Fe in $K_3[Fe(CN)_6]$ is

- A +2
- B +3
- C +1
- D +4

ATDB.uno

Question



In which of the following compound, iodine is in its highest oxidation state

- A KI
- B KIO_4
- C KI_3
- D IF_5

ATDB.uno



H.W module

Perubal + Perarambhan

Mole Concept

ATDB.uno



THANK
ATDB.uno

YOU