

PRAAYAS

JEE 2026

ATDB.uno

Mathematics

Basic Maths

Lecture - 09

By - Ashish Agarwal Sir
(IIT Kanpur)



Topics *To be covered*



- A** Logarithm and its Properties
- B** Problem Practice

ATDB.uno





Homework Discussion

ATDB.uno

QUESTION

TAH 05



$$\text{Solve : } (x^2 + 3x + 1)(x^2 + 3x - 3) \geq 5$$

$$(t + 1)(t - 3) \geq 5$$

$$t^2 - 2t - 3 \geq 5$$

$$t^2 - 2t - 8 \geq 0$$

$$(t - 4)(t + 2) \geq 0$$

ATDB.uno

$$(x^2 + 3x - 4)(x^2 + 3x + 2) \geq 0$$

$$(x + 4)(x - 1)(x + 1)(x + 2) \geq 0$$

$$\begin{array}{ccccccc} & + & & - & & + & & - & & + \\ & | & & | & & | & & | & & | \\ \hline & -4 & & -2 & & -1 & & 1 & & \end{array}$$

$$x \in (-\infty, -4] \cup [-2, -1] \cup [1, \infty)$$

QUESTION



TAH 06

Find Exhaustive set of values of x satisfying :

(i) $x^3 - 3x^2 - x + 3 > 0$

(ii) $x^4 - 3x^3 - x + 3 < 0$ $\xrightarrow{x^3(x-3) - 1(x-3) < 0}$
 $(x^3 - 1)(x - 3) < 0$

(iii) $x^4 + 6x^3 + 6x^2 + 6x + 5 \leq 0$ $\xrightarrow{(x-1)(x^2+x+1)(x-3) < 0}$
 $\xrightarrow{D < 0, a > 0 \rightarrow \text{always +ve}}$

$(x-1)(x-3) < 0$
 $x \in (1, 3)$

$x^3(x+1) + 5x^2(x+1) + x(x+1) + 5(x+1) \leq 0$

$(x^3 + 5x^2 + x + 5)(x+1) \leq 0$

$(x^2(x+5) + 1(x+5))(x+1) \leq 0$

$(x^2+1)(x+5)(x+1) \leq 0$

always +ve

$(x+5)(x+1) \leq 0 \xrightarrow{x \in [5, -1]}$

QUESTION

TAH 10




*$\Delta < 0$ always +ve
 $a > 0$*

Solve:
$$\frac{(x^2 - 4x + 5)^2 (x - 3)^2 (x + 1)^3}{(x - 1)(x - 5)^3 (x^2 - 7x + 12)} > 0$$

$$\frac{(x - 3)^2 (x + 1)^3}{(x - 1)(x - 5)^3 (x - 4)(x - 3)} > 0 \quad x \neq 3$$

$$\frac{(x - 3) (x + 1)^3}{(x - 1)(x - 4)(x - 5)^3}$$



$$x \in (-1, 1) \cup (3, 4) \cup (5, \infty)$$

ATDB.uno



Aao Machaay Dhamaal Deh Swaal pe Deh Swaal

ATDB.uno

QUESTION



Solve: $x(2^x - 1)(3^x - 9)(x - 3) < 0$.



$$x \cdot (x-0) \cdot (x-2) \cdot (x-3) < 0$$



$$x \in (2, 3)$$


$x=0$ is N.P

$$2^x - 1 = 0 \Rightarrow 2^x = 1 \Rightarrow x = 0$$



ATDB.uno

$$3^x - 9 = 0 \Rightarrow 3^x = 9 \Rightarrow x = 2$$



Ans. $x \in (2, 3)$


QUESTION



If $\frac{(e^x-1)(2x-3)(x^2+x+2)}{(\sin x-2)(x+1)x} \leq 0$ then $x \in$

D < 0, a > 0 always +ve.

-ve



~~A~~ $(-\infty, -1)$

~~B~~ $[\frac{3}{2}, \infty)$


C $(-1, 0)$

D $(0, \frac{3}{2})$

$\therefore -1 \leq \sin x \leq 1$
 $-3 \leq \sin x - 2 \leq -1$
 $\sin x - 2 \in \mathbb{R}^- \forall x \in \mathbb{R}$

$\frac{(e^x-1)(2x-3)}{x(x+1)} \geq 0$ *ATDB.uno* $\frac{x(2x-3)}{x(x+1)} \geq 0$ $e^x - 1 = 0 \Rightarrow x = 0$

$\frac{2x-3}{x+1} \geq 0, x \neq 0$



$e^x - 1$ behaves like $(x - 0)$

Ans. A, B

QUESTION



$$\text{Solve: } (x^2 - x - 1)(x^2 - x - 7) < -5$$

Tahoi

ATDB.uno

$$\text{Ans. } x \in (-2, -1) \cup (2, 3)$$

QUESTION



If $\frac{x^3(x-1)^2(x+4)}{(x+1)(x-3)} \geq 0$, then $x \in$

Tah02

A $(-\infty, -4]$

B $(-1, 0]$

C $(3, \infty)$

D $\{1\}$

ATDB.uno

Ans. A, B, C, D

QUESTION

TAH 03



1. Solve $\frac{x(3-4x)(x+1)}{(2x-5)} < 0$

[Ans. $x \in (-\infty, -1) \cup (0, 3/4) \cup (5/2, \infty)$]

2. Solve $\frac{(2x+3)(4-3x)^3(x-4)}{(x-2)^2x^5} \leq 0$


[Ans. $x \in (-\infty, -3/2) \cup (0, 4/3] \cup [4, \infty)$]

3. Solve $\frac{(x-3)(x+5)(x-7)}{|x-4|(x+6)} \leq 0$

$\frac{(x-3)(x+5)(x-7)}{(x+6)} \leq 0, x \neq 4$

[Ans. $x \in (-6, -5] \cup [3, 4) \cup (4, 7]$]

4. Solve $\frac{5x+1}{(x+1)^2} < 1$



$x \in (-6, -5] \cup [3, 7] - \{4\}$

OR

$x \in (-6, -5] \cup [3, 4) \cup (4, 7]$

[Ans. $x < 0$ or $x > 3, x \neq -1$]

5. Solve $\frac{x^4}{(x-2)^2} > 0$

[Ans. $x \in \mathbb{R} - \{0, 2\}$]

6. Solve $\frac{6x^2-5x-3}{x^2-2x+6} \leq 4$

[Ans. $-\frac{9}{2} \leq x \leq 3$]

7. Solve $\frac{(x+2)(x^2-2x+1)}{-4+3x-x^2} \geq 0$

[Ans. $x \in (-\infty, -2] \cup \{1\}$]

ATDB.uno



Logarithm



was invented to make calculations easier

logarithm converts product in to sum.
& converts division in to difference

ATDB.uno



Which would you Prefer to Attempt

(a) (37.53×8.74) or $37.53 + 8.74$

(b) $56.98 \div 12.76$ or $56.98 - 12.76$

ATDB.uno



Definition

$$\text{if } a^x = N \iff \log_a N = x$$

$(a > 0, a \neq 1, N > 0, x \in \mathbb{R})$

Input of logarithm (pointing to N)
 value of logarithm (pointing to x)
 Base of logarithm (pointing to a)

Ex: $\log_2 8 = 3$

Ex: $\log_2 x = -1$

find x : $2^{-1} = x$

$x = \frac{1}{2}$

$\log_a N$ is defined only if $a > 0, a \neq 1, N > 0$

ATDB.uno

* $\log_a N$ kaa value woh power hoti hai jisay 'a' pe lagaya jayay Taaki 'N' aa jayay

* $\log_a N = x \iff a^x = N$



JOHN NAPIER



Evaluate: $\log_{32} 128$

let $x = \log_{32} 128$

$$128 = 32^x$$

$$2^7 = 2^{5x}$$

$$5x = 7$$

$$x = 7/5$$

* $\log_a a = 1$

* $\log_a 1 = 0$

* $\log_a (1/a) = -1$

ATDB.uno

Evaluate: $\log_{10} (.01)$

$$\log_{10} \frac{1}{100}$$

"

$$\log_{10} 10^{-2}$$

"

$$-2$$

let $\log_{10} (0.01) = x$

$$0.01 = 10^x$$

$$\frac{1}{100} = 10^x$$

$$10^x = 10^{-2}$$

$$x = -2$$

* $\log_a \frac{1}{a} = -1$



$$a^{(\log_a x)} = x$$

proof:

$$\text{let } \log_a x = t \Rightarrow a^t = x$$

Now taking LHS

$$a^{\log_a x} = a^t = x = \text{RHS (proved)}$$

ATDB.uno

$$\text{Ex: } 2^{\log_2 100} = 100$$



Nichod!!



(a) $\log_N N = 1$ i.e. logarithm of a number to the same base is 1.

(b) $\log_{\frac{1}{N}} N = -1$ i.e. logarithm of a number of its reciprocal -1.

(c) $\log_a 1 = 0$ i.e. logarithm of unity to any base is zero.

(d) $a^{\log_a N} = N$ is a identify for all $N > 0$ and $a > 0, a \neq 1$

e.g. $2^{\log_2 5} = 5$

ATDB.uno

QUESTION



Column 1	Column 2
(a) $\log_{16} 32$ (S)	(P) -1
(b) $\log_9 27$ (T)	(Q) 1
(c) $\log_2(\log_2 4) = \log_2 2 = 1$ (Q)	(R) 2
(d) $\log_{2-\sqrt{3}}(2+\sqrt{3}) = -1$ (P)	(S) $5/4$
(e) $\log_{5\sqrt{5}} 125 = 2$ (R)	(T) $3/2$

$$\begin{aligned} \textcircled{a} \log_{16} 32 &= x \\ 32 &= 16^x \\ 2^5 &= 2^{4x} \\ x &= 5/4 \rightarrow \textcircled{S} \end{aligned}$$

$$\begin{aligned} \textcircled{b} \log_9 27 &= x \\ 9^x &= 27 \\ 3^{2x} &= 3^3 \\ x &= 3/2 \rightarrow \textcircled{T} \end{aligned}$$

ATDB.uno

$$(2+\sqrt{3})(2-\sqrt{3})=1$$

$2+\sqrt{3}$ & $2-\sqrt{3}$ are reciprocal of each other.



$$\log_2 4 = 2, \log_3 27 = 3, \log_{\frac{1}{2}} \frac{1}{8} = 3, \log_{\frac{1}{3}} \frac{1}{9} = 2$$


$\underbrace{\hspace{10em}}_{a, N > 1}$
 $\underbrace{\hspace{10em}}_{0 < a, N < 1}$

$$\log_2 \frac{1}{4} = -2, \log_{\frac{1}{3}} 27 = 3, \log_{\frac{1}{2}} 8 = -3$$

\downarrow
 \downarrow
 \downarrow

$a > 1, 0 < N < 1$
 $0 < a < 1, N > 1$

ATDB.uno



$$\log_a x = \log_a y \Rightarrow x = y$$

$$\log_a x = \log_a y = \lambda$$

$$x = a^\lambda = y$$



★ If $x=y \Rightarrow \log_a^x = \log_a^y$ (False)

★ If $x, y \in \mathbb{R}^+$ & $x=y$ then $\log_a^x = \log_a^y$ ($a > 0, a \neq 1$) (True)

ATDB.uno



NOTE:

1. It must be noted that whenever the number and the base are on the same side of unity the value of logarithm is positive, however if the number and the base are located on different side of unity then the value of logarithm is negative.
2. If two number are equal then their logarithm to the same base are equal and conversely.

ATDB.uno

QUESTION



Find the value of x.

$$(1) \log_{(5-x)}(x^2 - 2x + 65) = 2 \quad \leftarrow \quad (5-x)^2 = x^2 - 2x + 65$$

$$25 + x^2 - 10x = x^2 - 2x + 65$$

$$8x = -40$$

$$x = -5 \checkmark$$

$$(2) \log_{(x-1)}(4) = 2$$

$$(x-1)^2 = 4$$

$$x-1 = -2, 2$$

$$x = -1, 3 \checkmark$$

Base becomes -ve.

ATDB.uno

QUESTION



Find the value of x.

$$(3) \log_3(3^x - 6) = x - 1 \rightarrow 3^x - 6 = 3^{x-1}$$

$$3^x - 6 = 3^x \cdot 3^{-1}$$

$$(4) \log_2(4 + \log_3 x) = 3$$

$$3^x - 6 = \frac{3^x}{3}$$

$$\text{let } 3^x = t$$

$$4 + \log_3 x = 2^3$$

$$\log_3 x = 8 - 4 = 4$$

$$x = 3^4$$

$$x = 81$$

$$3t - 18 = t$$

$$3t - 18 = t$$

$$t = 9$$

$$3^x = 9$$

$$x = 2$$

ATDB.uno

QUESTION



Find the value of x.

$$(5) \log_2(x + 1) - \log_2(2x - 3) = 0$$

$$\log_2(x+1) = \log_2(2x-3)$$

$$x+1 = 2x-3$$

$$x=4$$

ATDB.uno

QUESTION



Find all values of x for which the following equalities hold true.

(i) $\log_2 x^2 = 1$

(ii) $\log_3 x = \log_3 (2 - x)$

(iii) $\log_4 x^2 = \log_4 x$

(iv) $\log_{1/2} (2x + 1) = \log_{1/2} (x + 1)$

(v) $\log_{1/3} (x^2 + 8) = -2$

Tah 04

ATDB.uno

QUESTION



$$\text{Solve: } 7^{\log_7 x} + 2x + 9 = 0$$

$$x + 2x + 9 = 0$$

$$x = -3 \text{ Ans}$$

correct: $x \in \phi$

Gadho / Gadhiyoo aigaa
naa kaoo

ATDB.uno

\therefore at $x = -3$ $\log_7 x$ is not defined.



The Principal Properties of Logarithm



If m, n are arbitrary positive numbers where, $a > 0, a \neq 1$ and x is any real numbers,

(1) $\log_a mn = \log_a m + \log_a n$ **proof:** let $\log_a m = x, \log_a n = y$

(2) $\log_a \frac{m}{n} = \log_a m - \log_a n$

(3) $\log_a m^x = x \log_a m$

ATDB.uno

proof: let $\log_a m = \alpha$
 $a^\alpha = m$
 $a^{\alpha x} = m^x$
 $\log_a m^x = \alpha x$
 $\log_a m^x = x \alpha = x \log_a m$

$a^x = m$ $a^y = n$
 $a^{x+y} = m \cdot n$ $\frac{a^x}{a^y} = \frac{m}{n}$
 $\log_a mn = x+y$ $a^{x-y} = m/n$
 $\log_a mn = \log_a m + \log_a n$
 $\log_a (m/n) = x-y$
 $\log_a m/n = \log_a m - \log_a n$

QUESTION



Find the value of following

$$(1) \log_{39} \frac{15}{7} + \log_{39} \frac{13}{3} - \log_{39} \frac{5}{21} \quad \leftarrow \log_{39} \left(\frac{15}{7} \cdot \frac{13}{3} \div \frac{5}{21} \right) = \log_{39} \left(\frac{15}{7} \times \frac{13}{3} \cdot \frac{21}{5} \right) = \log_{39} 39 = 1$$

$$(2) 2\log_6 2 + 3\log_6 3 + \log_6 12$$

$$\parallel$$

$$\log_6 2^2 + \log_6 3^3 + \log_6 12$$

ATDB.uno

$$\parallel$$

$$\log_6 (2^2 \cdot 3^3 \cdot 12) = \log_6 (2^2 \cdot 3^3 \cdot 2^2 \cdot 3^1)$$

$$= \log_6 (2^4 \cdot 3^4)$$

$$= \log_6 (2 \times 3)^4 = 4 \log_6 6 = 4$$

QUESTION



Find: $\log_2[\log_2\{\log_3(\log_3 27^3)\}]$

$$\log_2(\log_2(\log_3(\log_3 3^9)))$$

$$\log_2(\log_2(\log_3 9))$$

$$\log_2(\log_2 2)$$

$$\log_2 1 = 0$$

ATDB.uno

QUESTION



Solve:

(i) $2^{\log_2 x^2} - 3x - 4 = 0;$

(ii) $2^{2\log_2 x} - 3x - 4 = 0$

$$x^2 - 3x - 4 = 0 \quad (\because a^{\log_a x} = x)$$

$$(x-4)(x+1) = 0$$

$$x = 4, -1$$

$$2^{\log_2 x^2} - 3x - 4 = 0$$

$$x^2 - 3x - 4 = 0$$

$$x = 4, -1$$

ATDB.uno

$$\log_2 16 = \log_2 2^4 = 4 \log_2 2 = 4$$

$$\log_2 (-2)^4 \neq 4 \log_2 (-2)$$

$$4 \log_2 |-2| = 4 \cdot \log_2 2 = 4$$



$$\log_a x^{2n} = 2n \log_a |x|$$

(a) $\log_a mn$ can be written as $\log_a m + \log_a n$ (False)

ATDB.uno

(b) $\log_a m + \log_a n$ can be written as $\log_a mn$ (True)

$$\log_2(2 \cdot (-4)) \neq \log_2(-2) + \log_2(-4)$$



* find Domain of $f(x) = \log_{10}(x-1) + \log_{10}(x-4)$ $x-1 > 0$ & $x-4 > 0$

\checkmark
 $x > 4$

* find Domain of $g(x) = \log_{10}((x-1) \cdot (x-4))$ $D_f: (4, \infty)$

$\log_a N$ is defined only
if $a > 0, a \neq 1$ & $N > 0$

ATDB.uno

Domain: $x \in (-\infty, 1) \cup (4, \infty)$

QUESTION



If a , b and c are positive real numbers such that

$$a^{\log_3 7} = 27; b^{\log_7 11} = 49 \text{ and } c^{\log_{11} 25} = \sqrt{11}.$$

Find the value of $(a^{\log_3 7})^2 + b^{\log_7 11} + c^{\log_{11} 25}$.

$$E = (a^{\log_3 7})^{\log_3 7} + (b^{\log_7 11})^{\log_7 11} + (c^{\log_{11} 25})^{\log_{11} 25}$$

$$= 27^{\log_3 7} + 49^{\log_7 11} + \sqrt{11}^{\log_{11} 25}$$

$$= 7^{\log_3 27} + 11^{\log_7 49} + 25^{\log_{11} \sqrt{11}}$$

$$= 7^3 + 11^2 + 25^{1/2} = 469.$$

$$\therefore a^{\log_c b} = c^{\log_a b}$$



Chamatkaari BABA naa Banay.....



- ❖ $\log_a (m + n) \neq \log_a m + \log_a n$
- ❖ $\log_a (m + n) \neq \log_a m \cdot \log_a n$
- ❖ $\log_3 (9x) \neq 2 \log_3 3x$
- ❖ $\log_a^n x \neq n \log_a x$

$$\log_a^2 x = (\log_a x)^2$$

$$\log_a^3 x = (\log_a x)^3$$

$$\text{ATDB.uno } \log_a^n x = (\log_a x)^n$$



Base Changing Theorem



$$\frac{\log_c a}{\log_c b} = \log_b a \rightarrow \text{proof:}$$

$$\text{let } \log_b a = \alpha$$

$$a = b^\alpha$$

ATDB.uno

$$\begin{aligned} \text{LHS} \quad \frac{\log_c a}{\log_c b} &= \frac{\log_c b^\alpha}{\log_c b} = \frac{\alpha \log_c b}{\log_c b} = \alpha = \log_b a = \text{RHS} \end{aligned}$$

$$\log_b a = \frac{\log_c a}{\log_c b}$$

Ex: $\log_{32} 64$

$$\begin{aligned} &= \frac{\log_2 64}{\log_2 32} \\ &= \frac{\log_2 2^6}{\log_2 2^5} = \frac{6 \log_2 2}{5 \log_2 2} \\ &= \frac{6}{5} \end{aligned}$$



Important deduction from base changing theorem



$D_1: \log_b a = \frac{1}{\log_a b}$

proof: $\log_b a = \frac{\log_a a}{\log_a b} = \frac{1}{\log_a b}$

 $(a, b > 0, a, b \neq 1)$

$D_2: a^{\log_b c} = c^{\log_b a}$

Interchangeable

 $(a, b, c > 0, b \neq 1)$

proof:

$$a^{\log_b c} = a^{\frac{\log_a c}{\log_a b}} = \left(a^{\log_a c} \right)^{\frac{1}{\log_a b}} = c^{\log_a a} = c^{\log_b a}$$

Saari Class Illustrations ATDB.uno Retry karni Hai

QUESTION

(Home Challenge-02)



Solve in real numbers the equation

$$\sqrt{x_1 - 1} + 2\sqrt{x_2 - 4} + \dots + n\sqrt{x_n - n^2} = \frac{1}{2}(x_1 + x_2 + \dots + x_n) \text{ for } x_1, x_2, x_3, \dots, x_{n-1}, x_n.$$

ATDB.uno



Solution to Previous TAH

ATDB.uno

QUESTION**TAH 01**

$$\text{Solve: } (x - 1) (x^2 + 4x + 1) (x + 2) \leq 0$$

ATDB.uno

Sol

$$(x-1)(x^2+4x+1)(x+2) \leq 0$$


$$\hookrightarrow D = 16 - 4 = 12 > 0$$

$$\alpha, \beta = \frac{-4 \pm \sqrt{12}}{2} = -2 \pm \sqrt{3}$$

Tah-01

$$(x-1)(x - (-2 + \sqrt{3}))(x - (-2 - \sqrt{3}))(x+2) \leq 0$$

$$(x-1)(x+2-\sqrt{3})(x+2+\sqrt{3})(x+2) \leq 0$$



$$x \in [-(2+\sqrt{3}), -2] \cup [-(2-\sqrt{3}), 1] \quad \underline{\text{Ans}}$$



Date. / /

TAH-01

$$(x-1)(x^2+4x+1)(x+2) \leq 0$$

$$\rightarrow D = 16 - 4 \times 1 \times 1$$


$$D > 0$$

$$\alpha, \beta = \frac{-4 \pm \sqrt{12}}{2}$$

$$\alpha, \beta = -2 \pm \sqrt{3}$$

$$\Rightarrow (x-1)(x-(-2+\sqrt{3}))(x-(-2-\sqrt{3}))(x+2) \leq 0$$

-ve



$$x \in [-2-\sqrt{3}, -2] \cup [-2+\sqrt{3}, 1] \quad \text{Ans}$$

QUESTION

$$\text{Solve: } (x^2 - x - 6)(x^2 + 6x) \geq 0$$


TAH 02**ATDB.uno**



Tah 2 $\rightarrow (x^2 - x - 6)(x^2 + 6x) \geq 0$

$$(x-3)(x+2)x(x+6) \geq 0$$

ATDB.uno



$$(-\infty, -6] \cup [-2, 0] \cup [3, \infty)$$




$$\underline{\underline{Q}} \quad (x^2 - x - 6) (x^2 + 6x) \geq 0$$

Tah-02

Soln

$$(x^2 - 3x + 2x - 6) (x(x+6)) \geq 0$$

$$(x-3) (x+2) (x) (x+6) \geq 0 \rightarrow +ve$$



$$x \in (-\infty, -6] \cup [-2, 0] \cup [3, \infty) \quad \underline{\underline{\text{Ans}}}$$

QUESTION**TAH 03**

Solve: $x^2 - 5x + 6 \geq 0$ and $x^2 - 10x + 24 \leq 0$

ATDB.uno



110

Solve:

$$x^2 - 5x + 6 \geq 0$$

$$x^2 - 3x - 2x + 6 \geq 0$$

$$x(x-3) - 2(x-3) \geq 0$$

$$(x-2)(x-3) \geq 0$$



and $x^2 - 10x + 24 \leq 0$

$$x^2 - 6x - 4x + 24 \leq 0$$

$$x(x-6) - 4(x-6) \leq 0$$

$$(x-4)(x-6) \leq 0$$

Tah-03



ATDB.uno




$x \in [4, 6]$ Ans

Q $x^2 - 5x + 6 \geq 0$ and $x^2 - 10x + 24 \leq 0$

$$\Rightarrow x^2 - 3x - 2x + 6 \geq 0$$


$$\Rightarrow (x-3)(x-2) \geq 0$$



$$\Downarrow$$

$$x^2 - 6x - 4x + 24 \leq 0$$

$$(x-6)(x-4) \leq 0$$



$$x \in [4, 6]$$

Ans

Tah-03



QUESTION**TAH 04**

$$\frac{x^2 - x - 6}{x^2 + 6x} \geq 0$$

ATDB.uno



18

$$\frac{x^2 - x - 6}{x^2 + 6x} \geq 0$$

Tah - 04


SOM

$$\frac{x^2 - 3x + 2x - 6}{x(x + 6)}$$

ATDB.uno

$$\frac{(x-3)(x-2)}{x(x+6)} \geq 0$$

↳ +ve



$$x \in (-\infty, 6) \cup (0, 2] \cup [3, \infty)$$

Ans




Sol

$$\frac{x^2 - x - 6}{x^2 + 6x} \geq 0$$

$$\frac{x^2 - 3x + 2x - 6}{x(x+6)} = \frac{(x-3)(x+2)}{x(x+6)} \geq 0$$

Tahoy



$$x \in (-\infty, 6) \cup [-2, 0) \cup [3, \infty) \quad \underline{\text{Ans}}$$

QUESTION**TAH 05**

$$\text{Solve : } (x^2 + 3x + 1) (x^2 + 3x - 3) \geq 5$$

ATDB.uno



Q-5! Solve! $(x^2 + 3x + 1) \geq 5$

Soln!

Let $x^2 + 3x = t$.

$$\text{So, } (t+1)(t-3) \geq 5$$

$$\Rightarrow t^2 - 2t - 3 - 5 \geq 0$$

$$\Rightarrow t^2 - 2t - 8 \geq 0$$

$$\Rightarrow (t-4)(t+2) \geq 0$$

↓

$$\therefore t \in (-\infty, -2] \cup [4, \infty)$$

$$t \leq -2$$

or

$$t \geq 4$$

$$\text{or, } x^2 + 3x \leq -2$$

$$\Rightarrow x^2 + 3x + 2 \leq 0$$

$$\Rightarrow (x+1)(x+2) \leq 0$$

↓

$$x \in [-2, -1]$$

$$\text{or, } x^2 + 3x \geq 4$$

$$\text{or, } x^2 + 3x - 4 \geq 0$$

$$\text{or, } (x+4)(x-1) \geq 0$$

↓

$$x \in (-\infty, -4] \cup [1, \infty)$$

Union

$$\therefore x \in (-\infty, -4] \cup [-2, -1] \cup [1, \infty)$$

TAH 5
BY REED
FROM WB

ATDB.uno



$$(x^2 + 3x + 1)(x^2 + 3x - 3) \geq 5$$

Tah or

Sol

$$\text{Let, } x^2 + 3x = t$$

$$(t+1)(t-3) \geq 5$$

$$t^2 - 3t + t - 3 \geq 5$$

$$t^2 - 2t - 8 \geq 0$$


$$(t-4)(t+2) \geq 0$$

$$(x^2 + 3x - 4)(x^2 + 3x + 2) \geq 0$$

$$(x^2 + 4x - x - 4)(x^2 + 2x + x + 2) \geq 0$$

$$x(x+4) - 1(x+4) \quad x(x+2) + 1(x+2) \geq 0$$

$$(x-1)(x+4) \quad (x+1)(x+2) \geq 0$$



$$x \in (-\infty, -4] \cup [-2, -1] \cup [1, \infty)$$

QUESTION**TAH 06**

Find Exhaustive set of values of x satisfying :

(i) $x^3 - 3x^2 - x + 3 > 0$

(ii) $x^4 - 3x^3 - x + 3 < 0$

(iii) $x^4 + 6x^3 + 6x^2 + 6x + 5 \leq 0$

ATDB.uno



TAH 6 :- (ii) $x^4 - 3x^3 - x + 3 < 0$

$$x^3(x-3) - 1(x-3) < 0$$

$$\Rightarrow (x-3)(x^3-1) < 0$$

$$\Rightarrow (x-3)(x-1)(x^2+x+1) < 0$$

$D < 0$, $a > 1$ always +ve

$$\Rightarrow (x-3)(x-1) < 0$$

$$x \in (1, 3)$$

(iii) $x^4 + 6x^3 + 6x^2 + 6x + 5 \leq 0$

by hit & trial we get $(x+1)$ is factor

$$x^3(x+1) + 5x^2(x+1) + x(x+1) + 5(x+1) \leq 0$$

$$(x+1)(x^3 + 5x^2 + x + 5) \leq 0$$

$$(x+1)(x^2(x+5) + 1(x+5)) \leq 0$$

$$\Rightarrow (x+1)(x+5)(x^2+1) \leq 0$$

$D < 0$, $a > 0$, always +ve

30/04/2025 03:3




$$(n+1)(n+5)(n^2+1) \leq 0$$

always true

$$(n+1)(n+5) \leq 0$$

$$n \in [-5, -1]$$

ATDB.uno





Q-6! (i) $x^4 - 3x^3 - x + 3 < 0$

(ii) $x^4 + 6x^3 + 6x^2 + 6x + 5 \leq 0$

TAH 6-(i)

Soln \rightarrow (i) $x^4 - 3x^3 - x + 3 < 0$

or $x^3(x-3) - 1(x-3) < 0$

or $(x-3)(x^3-1) < 0$

or $(x-3)(x-1)(x^2+x+1) < 0$

$\rightarrow D < 0, a > 0 \Rightarrow$ always -ve.

or $(x-3)(x-1) < 0$

$\therefore x \in (1, 3)$



\rightarrow (ii) $x^4 + 6x^3 + 6x^2 + 6x + 5 \leq 0$

$\Rightarrow x^2(x^2 + 6x + 5) + (x^2 + 6x + 5) \leq 0$

$\Rightarrow (x^2 + 6x + 5)(x^2 + 1) \leq 0$

\downarrow
always +ve

$\Rightarrow (x+5)(x+1) \leq 0$

$\therefore x \in [-5, -1]$

TAH 6-(ii)
BY REED
FROM WB

QUESTION**TAH 07**

Solve: $(x - 1)^2 (x + 1)^3 (x - 4) < 0$


ATDB.uno



Tah-07

$$(x-1)^2 (x+1)^3 (x-4) < 0$$

ATDB.uno



$$x \in (-1, 1) - \{1\}$$



Q-7! Solve: $(x-1)^2(x+1)^3(x-4) < 0$


TAH 7

Solns

$$(x-1)^2(x+1)^3(x-4) < 0$$

⇓

$$(x-1)^2(x+1)(x-4) < 0 \quad \Delta \quad x \neq 1.$$



$$\therefore x \in (-1, 1) \cup (1, 4) - \{1\}$$

⇓

$$x \in (-1, 1) \cup (1, 4)$$

already
not present

QUESTION

TAH 08



$$\text{Solve: } \frac{(x-1)^2(x+1)^3}{x^4(x-2)} \leq 0$$


ATDB.uno



$$\textcircled{8} \frac{(x-1)^2 (x+1)^3}{x^4 (x-2)} \leq 0$$

$$\frac{\cancel{(x-1)^2} (x+1)^3}{(x-2)} \leq 0 \quad \begin{array}{l} x = 1 \checkmark \\ x \neq 0 \end{array}$$

ATDB.uno



$$x \in [-1, 2) - \{0\}$$



Q-8! Solve: $\frac{(x-1)^2 (x+1)^3}{x^4 (x-2)} \leq 0$


TAH 8

Soln

$$\frac{(x-1)^2 (x+1)^3}{x^4 (x-2)} \leq 0$$

$$\Rightarrow \frac{(x+1)}{(x-2)} \leq 0 \quad \& \quad \textcircled{x=1} \text{ is also possible}$$

$x \neq 0$



$$\therefore x \in [-1, 2) \cup \{1\} - \{0\}$$

$$\Rightarrow x \in [-1, 0) \cup (0, 2). \quad \text{[Ans]}$$

QUESTION

TAH 09



Find the exhaustive solutions set of

$$\frac{(2x - 5)^{100}(x + 3)(2x + 1)^{101}}{(x^2 - 4)^{151}(3x - 4)^{197}} < 0$$

ATDB.uno

$$\text{Ans. } (-\infty, -3) \cup \left(-2, -\frac{1}{2}\right) \cup \left(\frac{4}{3}, 2\right)$$




9)
$$\frac{(2x-5)^{100} (x+3) (2x+1)^{101}}{(x^2-4)^{151} (3x-4)^{197}} < 0$$

$x \neq 5/2 = 2.5$

$$\frac{(x+3) (2x+1)^{101}}{(x-2)^{151} (x+2)^{151} (3x-4)^{197}} < 0$$

↪ not in answer

$$x \in (-\infty, -3) \cup (-2, -1/2) \cup (4/3, 2)$$



[Handwritten signature]

Teacher's Sign



Q-71 Find the exponential


$$\frac{(2x-5)^{100} \cdot (x+3) \cdot (2x+1)^{101}}{(x^2-4)^{151} \cdot (3x-4)^{1017}} < 0$$

TAH 9
by Reed
from WB

Soln

$$\frac{(x+3) \cdot (2x+1)}{(x^2-4) \cdot (3x-4)} < 0 \quad \& \quad x \neq \frac{5}{2}$$

$$\Rightarrow \frac{(x+3) \cdot (2x+1)}{(x-2) \cdot (x+2) \cdot (3x-4)} < 0$$



$$\therefore x \in (-\infty, -3) \cup (-2, -\frac{1}{2}) \cup (\frac{4}{3}, 2) - \left\{ \frac{5}{2} \right\}$$

Ans: $x \in (-\infty, -3) \cup (-2, -\frac{1}{2}) \cup (\frac{4}{3}, 2)$

2.5
already
not
included



Q Find the exhaustive solutions set of


$$\frac{(2x-5)^{100} (x+3) (2x+1)^{101}}{(x^2-4)^{151} (3x-4)^{197}} < 0$$

Tah - 09

$$x \neq \frac{5}{2}$$

Sol

$$\frac{(x+3) (2x+1)^{101}}{(x-2)^{151} (x+2)^{151} (3x-4)^{197}} < 0$$



$$x \in (-\infty, -3) \cup (-2, -\frac{1}{2}) \cup (\frac{4}{3}, 2)$$

Ans

ATDB.uno

QUESTION

TAH 10



$$\text{Solve: } \frac{(x^2 - 4x + 5)^2 (x - 3)^2 (x + 1)^3}{(x - 1)(x - 5)^3 (x^2 - 7x + 12)} > 0$$

ATDB.uno



Q-10! Solve: $\frac{(x^2 - 4x + 5)^2 (x-3)^2 (x+1)^3}{(x-1) (x-5)^3 (x^2 - 7x + 12)} > 0$

TAH 10


Solⁿ ⇒ $D < 0$
 $a > 0$
 ↓
 always +ve.

↓
 $\frac{(x^2 - 4x + 5)^2 (x-3)^2 (x+1)^3}{(x-1) (x-5)^3 (x^2 - 7x + 12)} > 0$

↳ $D > 0 \Rightarrow$ factorizable.

⇒ $\frac{(x+1)^3 (x-3)^2}{(x-1) (x-5)^3 (x-3) (x-4)} > 0$

but $x-3 \neq 0$
 $\Rightarrow x \neq 3$



Since $0 > 0$ is not possible

∴ $x \in (-1, 1) \cup (3, 4) \cup (5, \infty)$
 ↓
 (Answer.)



Sol


$$\frac{(x^2 - 4x + 5)^2 (x-3)^2 (x+1)^2}{(x-1)(x-5)^3(x^2-7x+12)} > 0$$

$(x^2 - 4x + 5)^2$
 $a > 0, D < 0$
 always +ve

$x \neq 3$

Tah-10

$$\frac{(x+1)^3}{(x-1)(x-5)^3(x-3)(x-4)} > 0$$



$x \in (-1, 1) \cup (3, 4) \cup (5, \infty)$ Ans

QUESTION

TAH 11



Find the exhaustive solutions set of $\frac{(x-4)(2x-5)^{27}(x^2-9)^{10}(x+4)^{93}}{(x^2-25)(x+3)^{91}(x^2+10)^5} > 0$.

ATDB.uno

Ans. $(-\infty, -5) \cup (-4, -3) \cup \left(\frac{5}{2}, 3\right) \cup (3, 4) \cup (5, \infty)$




TAH-11

Q. Find exhaustive soln. set of $\frac{(x-4)(2x-5)^{27}(x^2-9)^{10}(x+4)^{93}}{(x^2-25)(x+3)^{91}(x^2+10)^5} > 0$

$\Rightarrow \frac{(x-4)(2x-5)^{27}(x+3)^{10}(x-3)^{10}(x+4)^{93}}{(x+5)(x-5)(x+3)^{91}(x^2+10)^5} > 0$ **TAH-11 BY NEELAKSH THAKUR SUPAUL**

∵ a > 0, b < 0
+ve always

$\Rightarrow \frac{(x-4)(2x-5)^{27}(x+4)^{93}}{(x+5)(x-5)(x+3)^{81}} > 0$, $x \neq -3$, $x = 3$ is not possible as, it not satisfy inequality.



$\therefore x \in (-\infty, -5) \cup (-4, -3) \cup (5/2, 4) \cup (5, \infty) - \{3\}$

→ already excluded

or, $(-\infty, -5) \cup (-4, -3) \cup (5/2, 3) \cup (3, 4) \cup (5, \infty)$

Ans



• Q-11:- Find the exhaustive solⁿ set of:

$$\frac{(x-4)(2x-5)^{29}(x^2-9)^{10}(x+4)^{93}}{(x^2-25)(x+3)^{91}(x^2+10)^5} > 0.$$

TAH 11
by Reed
from WB

Solⁿ:-


$$\frac{(x-4)(2x-5)^{29}(x^2-9)^{10}(x+4)^{93}}{(x^2-25)(x+3)^{91}(x^2+10)^5} > 0$$

$$\Rightarrow \frac{(x-4)(2x-5)(x+3)^{10}(x-3)^{10}(x+4)}{(x+5)(x-5)(x+3)^{91}(x^2+10)^5} > 0$$

$$\Rightarrow \frac{(x-4)(2x-5)(x-3)^{10}(x+4)}{(x+5)(x-5)(x+3)^{91}} > 0$$

D < 0 ⇒ always +ve.
a > 0

ATDB.uno & x ≠ -3.
x ≠ 3.



∴ x ∈ (-∞, -5) ∪ (-4, -3) ∪ (-5/2, 4) ∪ (5, ∞) - {3}.

↳ already excluded.

$$\Rightarrow x \in (-\infty, -5) \cup (-4, -3) \cup \left(\frac{5}{2}, 3\right) \cup (3, 4) \cup (5, \infty)$$

Answer.



Find the exhaustive solⁿ set of

$$\frac{(x-4)(2x-5)^{27}(x^2-9)^{10}(x+4)^{93}}{(x^2-25)(x+3)^{91}(x^2+10)^5} > 0$$


solⁿ

$$\frac{(x-4)(2x-5)^{27}(x-3)^{10}(x+3)^{10}(x+4)^{93}}{(x+5)(x-5)(x+3)^{91}(x^2+10)^5} > 0$$

$$\frac{(x-4)(2x-5)^{27}(x-3)^{10}(x+4)^{93}}{(x+5)(x-5)(x+3)^{91}(x^2+10)^5} > 0$$

$D = 0 - 40$
 $D = -40$
 $D < 0 - 9 > 0$
 $x = 3$ is eliminated b/c $2+10 > 0 \forall x \in \mathbb{R}$

$x = 3$ is eliminated (+ve region)



$$x \in (-\infty, 5) \cup (-4, -3) \cup (5/2, 4) \cup (5, \infty) - \{3\}$$

OR

$$x \in (-\infty, 5) \cup (-4, -3) \cup (5/2, 3) \cup (3, 4) \cup (5, \infty)$$

ans

for 3 as a critical point sign will remain same and it is excluded as for $x=3$ $0 > 0$ is false

QUESTION**TAH 12**

Solve the system of in equations $2x - 1 > x + \frac{7-x}{3} > 2, x \in \mathbb{R}$

ATDB.uno



Q. Solve the system of inequations $2x - 1 > x + \frac{7-x}{3} > 2, x \in \mathbb{R}$.

$$\Rightarrow 2x - 1 > x + \frac{7-x}{3} > 2$$

$$\Rightarrow 6x - 3 > 3x + 7 - x > 6$$

$$\Rightarrow 6x - 3 > 2x + 7 > 6$$

$$\Rightarrow 6x - 3 - 7 > 2x > 6 - 7$$

$$\Rightarrow 6x - 10 > 2x > -1$$

$$\Rightarrow 3x - 5 > x > -\frac{1}{2}$$

$$\Rightarrow 3x - 5 > x \text{ or, } x > \frac{5}{2}$$

$$\Rightarrow 2x > 5 \cap x > -\frac{1}{2}$$

$$\Rightarrow x > \frac{5}{2} \cap x > -\frac{1}{2}$$

$\therefore x > \frac{5}{2}$ or, $x \in \left(\frac{5}{2}, \infty\right)$ Ans

TAH-12 BY
NEELAKSH THAKUR
SUPAUL



• Q-12!
TAH-12

Solve the system of inequ,

$$2x-1 > x + \frac{7-x}{3} > 2; \quad x \in \mathbb{R}.$$

Solⁿ ⇒

$$2x-1 > x + \frac{7-x}{3} > 2$$

$$\Rightarrow 6x-3 > \frac{(2x+7)3}{3} > 6.$$

$$\Rightarrow 6x-3 > 2x+7 > 6$$

$$\Rightarrow 6x-3-7 > 2x > 6-7$$

$$\Rightarrow 6x-10 > 2x > -1$$

$$\Rightarrow 3x-5 > x > -\frac{1}{2}$$

$$\Rightarrow -\frac{1}{2} < x < 3x-5.$$

$$\text{or, } \left(x > -\frac{1}{2} \right) \left\{ \begin{array}{l} 3x-5 > x \\ \text{or, } 2x > 5 \\ \text{or, } \left(x > \frac{5}{2} \right) \end{array} \right.$$

$$\therefore \Rightarrow \boxed{x > \frac{5}{2}} \quad \therefore x \in \left(\frac{5}{2}, \infty \right)$$

Ans.

TAH.12
by Reed
from WB

ATDB.uno

QUESTION



TAH 13

Solve following double inequalities :

(i) $-3 < \frac{2x-7}{5} \leq 8$

(ii) $x^2 + 2 \leq 3x < 2x^2 - 5$

(iii) $-2 < \frac{x-5}{2x+1} < 5$

ATDB.uno



#TAH-13

TAH-13(i), (ii) BY NEELAKSH THAKUR SUPAUL

Q. Solve the following inequalities:-

(i). $-3 < \frac{2x-7}{5} \leq 8$

(ii). $x^2+2 \leq 3x \leq 2x^2-5$

(iii). $-2 < \frac{x-5}{2x+1} < 5$

⇒ (i). $-3 < \frac{2x-7}{5} \leq 8$

⇒ $-15 < 2x-7 \leq 8 \times 5$

⇒ $-15+7 < 2x \leq 40+7$

⇒ $-8/2 < x \leq 47/2$

∴ $-4 < x \leq 47/2$


or, $x \in (-4, 47/2]$

(ii). $x^2+2 \leq 3x \leq 2x^2-5$

$x^2+2-3x \leq 0 \cap 2x^2-5-3x \geq 0$

$(x-1)(x-2) \leq 0 \cap (x+1)(2x-5) \geq 0$

$x \in [1, 2] \cap x \in (-\infty, -1] \cup [5/2, \infty)$





Q-13! Solve:

TAH 13
BY REED
FROM WB

$$(i) \quad -3 < \frac{2x-7}{5} \leq 8$$

$$\Rightarrow -15 < 2x-7 \leq 40$$

$$\Rightarrow -8 \leq 2x \leq 47$$

$$\Rightarrow -4 \leq x \leq \frac{47}{2} \quad \therefore x \in \left[-4, \frac{47}{2}\right]$$

$$(ii) \quad x^2+2 \leq 3x < 2x^2-5$$

$$x^2+2-3x \leq 0$$

$$\Rightarrow (x-2)(x-1) \leq 0$$


$$\Downarrow \\ x \in [1, 2]$$

$$2x^2-5-3x > 0$$

$$\Rightarrow 2x^2-3x-5 > 0$$

$$\Rightarrow (2x-5)(x+1) > 0$$

$$\Downarrow \\ x \in (-\infty, -1) \cup \left(\frac{5}{2}, \infty\right)$$



$$\therefore x \in \emptyset$$



TAH-13 (iii) BY NEELAKSH THAKUR SUPAUL

$$\Rightarrow -2 < \frac{x-5}{2x+1} < 5$$


$$\Rightarrow -2 < \frac{x-5}{2x+1} \cap \frac{x-5}{2x+1} < 5$$

$$\Rightarrow \frac{x-5+4x+2}{2x+1} > 0 \cap \frac{x-5-10x-5}{2x+1} < 0$$

$$\Rightarrow \frac{5x-3}{2x+1} > 0 \cap \frac{-9x-10}{2x+1} < 0$$

$$\Rightarrow \frac{5x-3}{2x+1} > 0 \cap \frac{9x+10}{2x+1} > 0$$

$$x \in (-\infty, -1/2) \cup (3/5, \infty) \cap x \in (-\infty, -10/9) \cup (-1/2, \infty)$$



$$\therefore x \in (-\infty, -10/9) \cup (3/5, \infty)$$

$$\textcircled{110} \quad -2 < \frac{x-5}{2x+1} < 5$$

$$\Rightarrow \frac{x-5}{2x+1} > -2$$

$$\Rightarrow \frac{x-5+4x+2}{2x+1} > 0$$

$$\Rightarrow \frac{5x-3}{2x+1} > 0$$

$$\Downarrow$$

$$x \in (-\infty, -\frac{1}{2}) \cup (\frac{3}{5}, \infty)$$

$$\&$$

$$\frac{x-5}{2x+1} < 5$$

$$\Rightarrow \frac{x-5-10x-5}{2x+1} < 0$$


$$\Rightarrow \frac{9x+10}{2x+1} > 0$$

$$\Downarrow$$

$$x \in (-\infty, -\frac{10}{9}) \cup (-\frac{1}{2}, \infty)$$

$$\therefore x \in (-\infty, -\frac{10}{9}) \cup (\frac{3}{5}, \infty)$$

Answer





THANK
ATDB.uno

YOU