

PRAAYAS

JEE 2026

ATDB.uno

Mathematics

Basic Maths

Lecture - 13

By – Ashish Agarwal Sir
(IIT Kanpur)



Topics *To be covered*



- A** Introduction to Modulus
- B** Problem Practice

ATDB.uno





Homework Discussion

ATDB.uno

QUESTION [JEE Mains 2023]



The number of integral solutions x of $\log_{\left(x+\frac{7}{2}\right)} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$ is: $\left(\frac{x-7}{2x-3}\right)^2 > 0$
 $x \neq 7$ - (A)
 $x \neq \frac{3}{2}$

- A** 8
- B** 7
- C** 5
- ~~**D** 6~~

Case (i) if $x + \frac{7}{2} > 1 \Rightarrow x > -5/2$

Case (ii) if $0 < x + \frac{7}{2} < 1$
 $-\frac{7}{2} < x < -5/2$

$$\left(\frac{x-7}{2x-3}\right)^2 \geq 1$$

$$\frac{(x-7)^2}{(2x-3)^2} - 1 \geq 0$$

$$\frac{(x-7)^2 - (2x-3)^2}{(2x-3)^2} \geq 0$$

$$\frac{(3x-10)(-x-4)}{(2x-3)^2} \geq 0$$

ATDB.uno

$$\frac{(x-7)^2}{(2x-3)^2} \leq 1$$

$$\frac{(3x-10)(-x-4)}{(2x-3)^2} \leq 0$$


$$(3x-10)(x+4) \geq 0 \quad x \neq 3/2$$

$$x \in (-\infty, -4] \cup [10/3, \infty)$$





$$\frac{(3x-10)(x+4)}{(2x-3)^2} \leq 0, x \neq \frac{3}{2}$$



$$x \in [-4, \frac{10}{3}] - \{\frac{3}{2}\}$$

⇓

$$x \in (-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\}$$

ATDB.uno

$x \in \phi$

$$(-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\} \text{ --- (B)}$$

FINAL ANS : $A \cap B = (-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\}$

Integral values of $x = -2, -1, 0, 1, 2, 3 \Rightarrow 6$ values.

QUESTION



Solve the following inequalities:

$$(a) \log(x^2 - 2x - 2) \leq 0$$

$$(b) \log_5(x^2 - 11x + 43) < 2$$

$$(c) 2 - \log_2(x^2 + 3x) \geq 0$$

$$(d) \log_{1.5} \frac{2x-8}{x-2} < 0$$

$$(e) \log_3 \frac{1+2x}{1+x} < 1$$

$$(f) \log_4 \frac{3x+2}{x} \leq 0.5$$

$$(g) \log_2 \frac{x^2-4x+2}{x+1} \leq 1$$

$$(h) \log_2 \left(\frac{4x-3}{4-3x} \right) > -\frac{1}{2}$$

Answers:

$$(a) [-1, 1 - \sqrt{3}) \cup (1 + \sqrt{3}, 3];$$

$$(b) (2, 9);$$

$$(c) [-4, -3) \cup (0, 1];$$

$$(d) (4, 6);$$

$$(e) (-\infty, -2) \cup (-1/2, \infty);$$

$$(f) [-2, -2/3);$$

$$(g) [0, 2 - \sqrt{2}) \cup (2 + \sqrt{2}, 6];$$

$$(h) \left(\frac{3}{4}, \frac{4}{3} \right)$$



$$\textcircled{H} \left(\log_2 \left(\frac{4x-3}{4-3x} \right) \right)^2 \geq -\frac{1}{2}$$

As $\log_2 \left(\frac{4x-3}{4-3x} \right)$ should be defined

$$\frac{4x-3}{4-3x} > 0$$

ATDB.uno

$$\frac{4x-3}{3x-4} < 0$$

$$\begin{array}{c} + \quad - \quad + \\ \hline \quad | \quad | \\ \quad 3/4 \quad 4/3 \end{array}$$

$$x \in (3/4, 4/3) \text{ Ans}$$



Home Challenge-04

JEE MAINS 10 Apr Morning SHIFT

2023



Let a, b, c be three distinct positive real numbers such that

$(2a)^{\log_e a} = (bc)^{\log_e b}$ and $b^{\log_e 2} = a^{\log_e c}$. Then, $6a + 5bc$ is equal to

$$\ln(2a)^{\ln a} = \ln(bc)^{\ln b}$$

$$\ln a \cdot (\ln 2 + \ln a) = \ln b \cdot \ln bc$$

$$\ln a (\ln 2 + \ln a) = \ln b (\ln b + \ln c)$$

$$\ln 2 \cdot \ln a + (\ln a)^2 = (\ln b)^2 + \frac{\ln b \cdot \ln 2 \cdot \ln b}{\ln a}$$

$$(\ln a)^2 - (\ln b)^2 = \frac{(\ln b)^2 \cdot \ln 2}{\ln a} - \ln 2 \cdot \ln a = \ln 2 \left(\frac{(\ln b)^2}{\ln a} - \ln a \right) = \frac{\ln 2 ((\ln b)^2 - (\ln a)^2)}{\ln a}$$

$$\ln b^{\ln 2} = \ln a^{\ln c}$$

$$\ln 2 \cdot \ln b = \ln c \cdot \ln a$$

$$\ln c = \frac{\ln 2 \cdot \ln b}{\ln a}$$

ATDB.uno



$$(\ln a)^2 - (\ln b)^2 = \ln 2 \left(\frac{(\ln b)^2 - (\ln a)^2}{\ln a} \right)$$

$$(\ln a)^2 - (\ln b)^2 - \ln 2 \cdot \frac{(\ln b)^2 - (\ln a)^2}{\ln a} = 0$$

$$[(\ln a)^2 - (\ln b)^2] \left(1 + \frac{\ln 2}{\ln a} \right) = 0$$

$$(\ln a)^2 - (\ln b)^2 = 0 \quad \text{or} \quad \frac{\ln 2}{\ln a} + 1 = 0$$

$$(\ln a - \ln b)(\ln a + \ln b) = 0$$

$$\ln a = \ln b \quad \text{or} \quad \ln a + \ln b = 0$$

$$a = b \quad \text{or} \quad \ln a = -\ln b = \ln \frac{1}{b}$$

(N.P) $a = \frac{1}{b}$

$$\ln 2 = -\ln a$$

$$\ln a = \ln \frac{1}{2}$$

$$a = \frac{1}{2} \Rightarrow \ln c = \frac{\ln 2 \cdot \ln b}{\ln a} = \frac{\ln 2 \cdot \ln b}{-\ln 2} = -\ln b \Rightarrow \ln b + \ln c = 0$$

$$\ln bc = 0 \Rightarrow bc = 1$$

~~$$x^2 = x$$~~

$$x = 1$$

Gadho/Gadhiyoo a isaa
naa kano!!

Sahi

$$x^2 = x$$

$$x^2 - x = 0 \Rightarrow x(x-1) = 0$$

$$x = 0, 1$$



$$\text{If } a = \frac{1}{b}$$

$$\ln c = \frac{\ln 2 \cdot \ln b}{\ln a}$$

$$\ln c = \frac{\ln 2 \cdot \ln b}{-\ln b} = -\ln 2$$

$$c = \frac{1}{2}$$

Hence

$$6a + 5bc = 6 \cdot \frac{1}{2} + 5 = \underline{8 \text{ Ans}}$$

ATDB.uno

$6a + 5bc$ has ∞ many possible values.

$$(a^2)^{\ln a} = (bc)^{\ln b}, \ln 2 = a^{\ln c} \rightarrow b^{\ln 2} = \left(\frac{1}{b}\right)^{\ln 1/2} = (b^{-1})^{-\ln 2} = b^{\ln 2}$$

$$\left(\frac{2}{b}\right)^{\ln \frac{1}{b}} = \left(\frac{b}{2}\right)^{\ln b}$$

$$\left(\frac{2}{b}\right)^{-\ln b} = (b/2)^{\ln b} \Rightarrow (b/2)^{\ln b} = (b/2)^{\ln b}$$



Aao Machaay Dhamaal Deh Swaal pe Deh Swaal

ATDB.uno

QUESTION [JEE Mains 2021]

The number of solutions of the equation $\log_4(x - 1) = \log_2(x - 3)$ is

Tahoi

ATDB.uno

QUESTION



Find x for $\frac{(\ln x)^2 - 3 \ln x + 3}{\ln x - 1} < 1.$

$$\ln x = t$$

$$\frac{t^2 - 3t + 3}{t - 1} - 1 < 0$$

$$\frac{t^2 - 4t + 4}{t - 1} < 0$$

$$\frac{(t - 2)^2}{(t - 1)} < 0$$

$$\frac{1}{t - 1} < 0 \quad t \neq 2$$

$$t - 1 < 0$$

$$t < 1$$

ATDB.uno

$$\ln x < 1 \quad \& \quad x > 0, \quad \ln x \neq 2$$

$$\log_e x < 1$$

$$x < e$$

$$x \neq e^2$$

$$x \in (0, e)$$

QUESTION



$$\text{Solve : } \log_{\frac{1}{x}} \left(\frac{2(x-2)}{(x+1)(x-5)} \right) \geq 1$$

Tah02

ATDB.uno




Modulus/Absolute value Function



$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0 \end{cases}$$

★ $|x| \geq 0$

★ $|x| = 0 \iff x = 0$




ATDB.uno



Geometrical Meaning of Modulus



$|x|$ = distance of x on number line from 0



Ex: $|x| = 2$

$x = -2, 2$

Ex: $|2x - 1| = 5$

$2x - 1 = -5, 5$

$2x = -4, 6$

$x = -2, 3$

Ex: $|2x - 6| = -10$

→ (No soln)

$|x| = a, a > 0$
 \Downarrow
 $x = \pm a$



Important Properties



$$P_1: |-x| = |x| \quad \star\star\star\star\star$$

$$P_2: \left| \frac{x}{y} \right| = \frac{|x|}{|y|}$$

Module distributes
over multiplication
& division

$$|x_1 \cdot x_2 \cdots x_n| = |x_1| \cdot |x_2| \cdots |x_n|$$

$$P_3: |xy| = |x||y|$$



Important Properties

P₄: $\sqrt{x^2} = |x|, x \in \mathbb{R}$, Also $x^2 = |x|^2$

Yaad Rakhe (variable/Expression)²
 ko $|variable/Expression|^2$
 likh sakte hai

P₅: $|x| \geq 0$ also $|x| \geq x$

✦ $|x| > x \rightarrow x \in \mathbb{R}^-$

✦ $|x| = x \rightarrow x \in [0, \infty)$

ATDB.uno



$$(i) |f(x)| = a, a \geq 0$$

$$f(x) = \pm a$$

$$(ii) |f(x)| = g(x)$$

$$\text{Case ① } f(x) \geq 0$$

$$\downarrow \text{ solve } f(x) = g(x)$$

$$\text{Case ② } f(x) < 0$$

$$-f(x) = g(x)$$

$$|f(x)| = |g(x)|$$

$$\downarrow$$

$$(f(x))^2 = (g(x))^2$$

$$\text{Apply } a^2 - b^2$$

$$\text{||}$$

$$(a-b)(a+b)$$

UNION

(FINAL Ans)


QUESTION

$$|x^2 + x - 20| = -(x^2 + x - 20)$$

$$\Downarrow$$
$$x^2 + x - 20 \leq 0$$

$$\Downarrow$$
$$(x+5)(x-4) \leq 0$$

$$x \in [-5, 4]$$



ATDB.uno



QUESTION



$$\left| \left(\frac{x^2 - 6x + 8}{x^2 - 4x + 3} \right) \right| = - \left(\frac{x^2 - 6x + 8}{x^2 - 4x + 3} \right)$$

Tah03

ATDB.uno

QUESTION



$$\text{Solve: } |2x - 1| = 7$$

$$2x - 1 = -7,7$$

$$2x = -6,8$$

$$x = -3,4$$

ATDB.uno

QUESTION



Solve: $|3 - x| = 2$

$3 - x = -2, 2$

$-x = -5, -1$

$x = 5, 1$

ATDB.uno

QUESTION



$|x + 2| = 2(3 - x)$ then x is equal to

~~A~~ $4/3$

B -8

C 8

D $-4/3$

Case ① if $x + 2 > 0 \Rightarrow x > -2$

$$x + 2 = 6 - 2x$$

$$3x = 4$$

$$x = \frac{4}{3}$$

Case ② if $x + 2 < 0 \Rightarrow x < -2$

$$-x - 2 = 6 - 2x$$

$$x = 8$$

$x \in \phi$

ATDB.uno

$$x = \frac{4}{3}$$

QUESTION



If $|3x - 2| + x = 11$ then x is

Tahoy

- A** $13/4$
- B** $9/2$
- C** $-9/2$
- D** $-13/4$

ATDB.uno

QUESTION



$||x - 2| - 2| - 2| = 2$ then sum of all values of x satisfying the equation is

~~A~~ 8

B 12

C 0

D 4

$$||x - 2| - 2| - 2| = 2$$

$$|m - 2| = 2$$

$$m - 2 = -2, 2$$

$$m = 0, 4$$

ATDB.uno

$$m = 0$$

$$|x - 2| - 2 = 0$$

$$|x - 2| - 2 = 0$$

$$|x - 2| = 2$$

$$x - 2 = -2, 2$$

$$x = 0, 4$$

$$m = 4$$

$$|x - 2| - 2 = 4$$

$$|x - 2| - 2 = -4, 4$$

$$|x - 2| = -2, 6$$

$$|x - 2| = -2 \text{ or } |x - 2| = 6$$

Not possible

$$x - 2 = 6, -6$$

$$x = 8, -4$$



$$x = 0, 4, 8, -4$$

$$\text{Sum} = 0 + 4 + 8 - 4 = 8$$

ATDB.uno

QUESTION



$$\text{Solve: } |x + 1| + |x - 2| = 5$$

F_1	-ve	+ve	+ve
F_2	-ve	-ve	+ve

Case I

if $x > 2$

$$x + 1 + x - 2 = 5$$

$$2x = 6$$

$$x = 3$$

$x = 3$

Case II $-1 < x < 2$

$$x + 1 - (x - 2) = 5$$

$$3 = 5$$

↓
Not possible

Case III if $x \leq -1$

$$-(x + 1) - (x - 2) = 5$$

$$-2x + 1 = 5$$

$$x = -2$$

$x = -2$

UNION

Ans $x = -2, 3$



QUESTION

$|x - 1| + |x - 2| + |x - 3| = 9$ then x can be

Tah 05

- A** -5
- B** 9
- C** -1
- D** 5

T_1	-ve	+	+	+
T_2	-ve	-ve	+	+
T_3	-ve	-ve	-ve	+

Case ① if $x \leq 1$

$$-(x-1) - (x-2) - (x-3) = 9$$

$$-3x + 6 = 9$$

$$x = -1$$

x = -1

Case ② if $1 < x \leq 2$

Case ③

ATDB.uno

QUESTION [JEE Mains 2021]



The number of real solution of the equation, $x^2 - |x| - 12 = 0$ is:

- A** 2
B 3
C 1
D 4

M(1)

case (i) if $x \geq 0$

$$x^2 - x - 12 = 0$$

$$(x-4)(x+3) = 0$$

$$x = -3, 4$$

case (ii) if $x < 0$

$$x^2 + x - 12 = 0$$

$$(x+4)(x-3) = 0$$

$$x = -4, 3$$

ATDB.uno

$$x = -4, 4$$

M(2)

$$x^2 - |x| - 12 = 0$$

$$|x|^2 - |x| - 12 = 0$$

$$|x| = t \quad t^2 - t - 12 = 0$$

$$t = 4, -3$$

~~$$|x| = 4, -3$$~~

$$x = \pm 4$$

QUESTION [JEE Mains 2025 (8 April)]

Tah 06



The sum of the squares of the roots of $|x - 2|^2 + |x - 2| - 2 = 0$ and the squares of the roots of $x^2 - 2|x - 3| - 5 = 0$, is

A 24

B 26

C 36

D 30

$$|x - 2| = t$$

$$t^2 + t - 2 = 0$$

$$(t + 2)(t - 1) = 0$$

$$t = -2, 1$$

ATDB.uno

~~$$|x - 2| = -2, 1$$~~

Ans. C

QUESTION [JEE Mains 2025 (7 April)]



The number of real roots of the equation $x|x - 2| + 3|x - 3| + 1 = 0$ is:

Tah 07

A 4

B 3

C 2

D 1

$$\begin{array}{cccc}
 T_1 & -ve & | & +ve & | & +ve \\
 T_2 & -ve & 2 & -ve & 3 & +ve
 \end{array}$$

case 1

ATDB.uno

Ans. D

QUESTION [JEE Mains 2024 (5 April)]

The number real solutions of the equations $x|x + 5| + 2|x + 7| - 2 = 0$ is

Tahos

ATDB.uno

Ans. 3

QUESTION [JEE Mains 2024 (5 April)]



The number of distinct real roots of the equation $|x| |x + 2| - 5|x + 1| - 1 = 0$ is

Tah 09

T_1	-ve	-ve	-ve	+
T_2	-ve - 2	+ve - 1	+ve 0	+
T_3	-ve	-ve	+ve	

Case ① if $x \leq -2$

$$-x \cdot -(x+2) + 5(x+1) - 1 = 0$$

$$x^2 + 2x + 5x + 5 - 1 = 0$$

$$x^2 + 7x + 4 = 0$$

$$x = \frac{-7 \pm \sqrt{49 - 16}}{2} = \frac{-7 \pm \sqrt{33}}{2}$$

$$\frac{-7 + \sqrt{33}}{2} > -2$$

$$\frac{-7 + \sqrt{33}}{2} > -4$$

$$\sqrt{33} > 3$$


$$x = \frac{-7 - \sqrt{33}}{2}$$

Ans. 3

QUESTION [JEE Mains 2024 (8 April)]



The number of distinct real roots of the equation $|x + 1| |x + 3| - 4|x + 2| + 5 = 0$ is



Case (i) if $x \leq -3$

Case (ii) if $-3 < x \leq -2$

Case (iii) if $-2 < x < -1$

Case (iv) if $x > -1$

ATDB.uno

Thumbs up
Thumbs down

Yes understood - 96%

Abhi bhi nahi aaya - 4%



No: of Number
 ke Tukde
 ||
 No: of cases

Ans. 2

QUESTION [JEE Mains 2023]



The sum of all the roots of the equation $|x^2 - 8x + 15| - 2x + 7 = 0$ is:

- A** $11 + \sqrt{3}$
- ~~**B** $9 + \sqrt{3}$~~
- C** $9 - \sqrt{3}$
- D** $11 - \sqrt{3}$

case ① If $x^2 - 8x + 15 > 0$

$$(x-3)(x-5) > 0$$

$$\begin{array}{c} + \quad - \quad + \\ | \quad | \quad | \\ 3 \quad 5 \end{array}$$

$$x \in (-\infty, 3) \cup (5, \infty)$$

case ②

If $x^2 - 8x + 15 \leq 0$
 $(x-5)(x-3) \leq 0$

$$x \in [3, 5]$$

$$x^2 - 8x + 15 - 2x + 7 = 0$$

$$x^2 - 10x + 22 = 0$$

$$x = \frac{10 \pm \sqrt{12}}{2}$$

$$x = 5 \pm \sqrt{3}$$

$$-(x^2 - 8x + 15) - 2x + 7 = 0$$

$$-x^2 + 6x - 8 = 0$$

$$x^2 - 6x + 8 = 0$$

$$x = 2, 4$$

ATDB.uno

\cap

$$x = 5 + \sqrt{3}$$

UNION

$$x = 4, 5 + \sqrt{3}$$

$$\text{sum} = 9 + \sqrt{3}$$

$$x = 4$$

\cap

QUESTION [JEE Mains 2024 (30 Jan)]



The number of real solutions of the equation $x(x^2 + 3|x| + 5|x - 1| + 6|x - 2|) = 0$ is _____

$$x = 0 \quad \text{or} \quad \underbrace{x^2}_{\geq 0} + \underbrace{3|x|}_{\geq 0} + \underbrace{5|x-1|}_{\geq 0} + \underbrace{6|x-2|}_{\geq 0} = 0$$

$x = 0$ (only one soln)
 $x^2 = 0 \Rightarrow x = 0$
 $x - 1 = 0 \Rightarrow x = 1$
 $x - 2 = 0 \Rightarrow x = 2$

$x = 0$
 $x = 0$
 $x = 1$
 $x = 2$

$x \in \phi$
 NO soln

Ans. 1



QUESTION [JEE Mains 2018]

Let $S = \{x \in \mathbb{R}: x \geq 0 \text{ and } 2|\sqrt{x} - 3| + \sqrt{x}(\sqrt{x} - 6) + 6 = 0\}$. Then S :

- ~~A~~ contains exactly two elements
- B contains exactly four elements
- C is an empty set
- D contains exactly one element

$$\sqrt{x} = t$$

$$2|t-3| + t(t-6) + 6 = 0$$

Case ① $t \geq 3$

$$2(t-6) + t^2 - 6t + 6 = 0$$

$$t^2 - 4t = 0$$

$$t = 0, 4$$

$$t = 4$$

Case ② if $t < 3$

$$-2t + 6 + t^2 - 6t + 6 = 0$$

$$t^2 - 8t + 12 = 0$$

$$t = 2, 6$$

$$t = 2$$

ATDB.uno

$$t = 4, 2$$

$$\sqrt{x} = 4, 2$$

$$S = \{16, 4\} \quad x = 16, 4$$

QUESTION [JEE Mains 2021]

Tah09

The number of the real roots of the equation $(x + 1)^2 + |x - 5| = \frac{27}{4}$ is

ATDB.uno



Sabse Important Baat



Sabhi Class Illustrations Retry Karnay hai...

ATDB.uno



Today's BPP

ATDB.uno



Lo Karo Duvaadaar Practice!!



1. $\log_5(x^2 - 3x + 3) > 0$

2. $\log_7[\log_5(x^2 - 7x + 15)] > 0$

3. $\log_{\left(\frac{1}{2}\right)}[\log_5(x^2 - 7x + 17)] > 0$

4. $\log_{\left(\frac{1}{2}\right)}(\log_5(\log_2(x^2 - 6x + 40))) > 0$

5. $\log_3[\log_5 \log_2(x^2 - 9x + 50)] > 0$

6. $\log_6\left(\frac{x-2}{6-x}\right) > 0$

7. $\log_{0.5}(x^2 - 5x + 6) > -1$

8. $\log_8(x^2 - 4x + 3) < 1$

9. $\log_{\left(\frac{1}{4}\right)}\left(\frac{35-x^2}{x}\right) \geq -\frac{1}{2}$



Answers



1. $(-\infty, 1) \cup (2, \infty)$

2. $(-\infty, 2) \cup (5, \infty)$

3. $(3, 4)$

4. $(2, 4)$

5. $(-\infty, 3) \cup (6, \infty)$

6. $(4, 6)$

7. $(1, 4)$

8. $(-1, 5)$

9. $(-1, 0) \cup (5, \infty)$

ATDB.uno



Home Challenge-05



If the value of x which satisfies the equation $2 \log_3 \sqrt{3^{1-x} + 2} = 1 + \log_3(4 \cdot 3^x - 1)$ is given by, $1 - \log_3 k$, then find the value of k . [Ans. 4]

ATDB.uno



Homework From Module



Prarambh (Topicwise) : Q1 to Q17

Prabal (JEE Main Level) : Q1 to Q7

ATDB.uno



Solution to Previous TAH

ATDB.uno

QUESTION [JEE Mains 2023]

The number of integral solutions x of $\log_{\left(x+\frac{7}{2}\right)} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$ is :

A 8

B 7

C 5

D 6

ATDB.uno


Q-3 (TAH-1): The number of integral solutions x of $\log_{x+\frac{7}{2}} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$ is:
 (A) 8 (B) 7 (C) 5 (D) 6.

TAH-1, By Reed from WB

Solⁿ: $\left(\frac{x-7}{2x-3}\right)^2 > 0$ for log to be defined
 $\Rightarrow \frac{x-7}{2x-3} \neq 0 \Rightarrow x \neq 7, x \neq \frac{3}{2}$ — (A)

Case-1: If $x + \frac{7}{2} > 1$
 $\Rightarrow x > 1 - \frac{7}{2} \Rightarrow x > -\frac{5}{2}$ — (P.C.)

$$\begin{aligned} \therefore \log_{x+\frac{7}{2}} \left(\frac{x-7}{2x-3}\right)^2 &\geq 0 \\ \Rightarrow \left(\frac{x-7}{2x-3}\right)^2 &\geq 1 \\ \Rightarrow \frac{(x-7)^2}{(2x-3)^2} - 1 &\geq 0 \\ \Rightarrow \frac{(x-7)^2 - (2x-3)^2}{(2x-3)^2} &\geq 0 \\ \Rightarrow \frac{(x-7+2x-3)(x-7-2x+3)}{(2x-3)^2} &\geq 0 \\ \Rightarrow \frac{(3x-10)(-x-4)}{(2x-3)^2} &\geq 0 \\ \Rightarrow (3x-10)(-x-4) &\leq 0, x \neq \frac{3}{2} \end{aligned}$$



Taking 'n' with P.C.
 $x > -\frac{5}{2} \cap x \in [-4, \frac{10}{3}] - \{\frac{3}{2}\}$
 $\therefore x \in (-\frac{5}{2}, \frac{3}{2}) \cup (\frac{3}{2}, \frac{10}{3}]$ — (B)


TAH 1(part-1) by Reed from WB

Case-2: If $0 < x + \frac{7}{2} < 1$.
 or, $-\frac{7}{2} < x < -\frac{5}{2}$ — (P.C.)


TAH 1(part 2) by Reed from WB

$$\begin{aligned} \text{now, } \log_{x+\frac{7}{2}} \left(\frac{x-7}{2x-3}\right)^2 &\geq 0 \\ \Rightarrow \left(\frac{x-7}{2x-3}\right)^2 &\leq 1 \\ \Rightarrow \frac{(3x-10)(x+4)}{(2x-3)^2} &\geq 0 \\ \Rightarrow (3x-10)(x+4) &\geq 0, x \neq \frac{3}{2} \end{aligned}$$

$\therefore x \in (-\infty, -4] \cup [\frac{10}{3}, \infty)$



$(\text{Case-1}) \cup (\text{Case-2}) \Rightarrow \text{Case-0}$
 $x \in (-\frac{5}{2}, \frac{3}{2}) \cup (\frac{3}{2}, \frac{10}{3}]$ — (B)



$\therefore x \in (-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\} \rightarrow$ final condition of 'x'.

\therefore Integral values in this region
 $= -2, -1, 0, 1, 2, 3$.
 6 integral values. \therefore Ans \Rightarrow (B)



QUESTION

$$\log_{x+3}(x^2 - x) < 1$$

ATDB.uno



Tah 2

$\log_{x+3} (x^2-x) < 1$

Also $x+3 > 0$
 $x > -3$
No Need

Also $x-x > 0$
 $x(x-1) > 0$

$x \in (-\infty, 0) \cup (1, \infty)$
 (1)

Case I: if $x+3 > 1 \Rightarrow x > -2$

Case II: $0 < x+3 < 1$

$x^2-x < x+3$

$x^2-2x-3 < 0$

$(x-3)(x+1) < 0$

$x \in (-1, 3)$

$x \in (-1, 3)$ - (2)

$x^2-x > x+3$

$x^2-2x-3 > 0$

$(x-3)(x+1) > 0$

$x \in (-\infty, -1) \cup (3, \infty)$


$-3 < x < -2$

$x \in (-3, -2)$ - (3)

Sakshi sahu

(2) \cup (3)

$x \in (-3, -2) \cup (-1, 3) \cap (1)$



$x \in (-3, -2) \cup (-1, 0) \cup (1, 3)$ sh



Kriti Mathur Raj.



TAH2) $\log_{x+3}(x^2-x) < 1$



Case 1: If $x+3 > 1$
 $x > -2$
 $x \in (-2, \infty)$ - (A)

$x^2-x < x+3$
 $x^2-x-x-3 < 0$
 $x^2-2x-3 < 0$
 $x^2-3x+x-3 < 0$
 $x(x-3)+1(x-3) < 0$
 $(x+1)(x-3) < 0$





Case 2: If $0 < x+3 < 1$
 $-3 < x < -2$
 $x \in (-3, -2)$ - (C)

$x^2-x > x+3$
 $x^2-x-x-3 > 0$
 $x^2-2x-3 > 0$
 $(x+1)(x-3) > 0$
 $x \in (-\infty, -1) \cup (3, \infty)$ - (D)



Now,

$x^2-x > 0$
 $x(x-1) > 0$



$x \in (-3, -2) \cup (-1, 0) \cup (1, 3)$ Ans.

ATDB.uno




TAH-2: $\log_{x+3}(x^2-x) < 1$.

Solve!

$$\log_{x+3}(x^2-x) < 1$$

for log to be defined,

$$\begin{matrix} x+3 > 0 & | & x+3 \neq 1 & | & x^2-x > 0 \\ \Rightarrow x > -3 & | & \Rightarrow x \neq -2 & | & \Rightarrow x(x-1) > 0 \\ & & & & \Rightarrow x \in (-\infty, 0) \cup (1, \infty) \end{matrix}$$



$\therefore x \in (-3, 0) \cup (1, \infty) - \{-2\}$

TAH 2
by Reed
From WB

Case-1: If $x+3 > 1 \Rightarrow x > -2$ — (P.C)

So, $\log_{x+3}(x^2-x) < 1$

$$\begin{aligned} \Rightarrow (x^2-x) &< x+3 \\ \Rightarrow x^2-2x-3 &< 0 \\ \Rightarrow (x-3)(x+1) &< 0 \end{aligned}$$



$\therefore x \in (-1, 3)$ — (P.C)

Case-2: If $x+3 \in (0, 1)$.

$$\begin{aligned} \Rightarrow 0 < x+3 < 1 \\ \Rightarrow -3 < x < -2 \end{aligned} \quad \therefore x \in (-3, -2) \text{ — (P.C)}$$

So, $\log_{x+3}(x^2-x) < 1$

$$\begin{aligned} \Rightarrow (x^2-x) &> (x+3) \\ \Rightarrow x^2-2x-3 &> 0 \\ \Rightarrow (x-3)(x+1) &> 0 \end{aligned}$$

$\therefore x \in (-\infty, -1) \cup (3, \infty)$




$\therefore x \in (-3, -2)$ — (P.C)

Taking (1) \cup (2) $\Rightarrow x \in (-3, -2) \cup (-1, 3)$ — (P.C)

Taking intersection of (1) & (2):

$A \cap B \Rightarrow$



$\therefore x \in (-3, -2) \cup (-1, 3)$



Solution to Previous BPPs

ATDB.uno

QUESTION



Solve the following inequalities:

$$(a) \log(x^2 - 2x - 2) \leq 0$$

$$(b) \log_5(x^2 - 11x + 43) < 2$$

$$(c) 2 - \log_2(x^2 + 3x) \geq 0$$

$$(d) \log_{1.5} \frac{2x-8}{x-2} < 0$$

$$(e) \log_3 \frac{1+2x}{1+x} < 1$$

$$(f) \log_4 \frac{3x+2}{x} \leq 0.5$$

$$(g) \log_2 \frac{x^2-4x+2}{x+1} \leq 1$$

$$(h) \log_2 \left(\frac{4x-3}{4-3x} \right) > -\frac{1}{2}$$

Answers:

$$(a) [-1, 1 - \sqrt{3}) \cup (1 + \sqrt{3}, 3];$$

$$(b) (2, 9);$$

$$(c) [-4, -3) \cup (0, 1];$$

$$(d) (4, 6);$$

$$(e) (-\infty, -2) \cup (-1/2, \infty);$$

$$(f) [-2, -2/3);$$

$$(g) [0, 2 - \sqrt{2}) \cup (2 + \sqrt{2}, 6];$$

$$(h) \left(\frac{3}{4}, \frac{4}{3} \right)$$



BPPs: Solve the following meq.s.

Solⁿ ⇒ **(a)** $\log(x^2 - 2x - 2) \leq 0$

⇒ $x^2 - 2x - 2 \leq 1$

⇒ $x^2 - 2x - 3 \leq 1$

⇒ $(x-3)(x+1) \leq 0$

⇒ $x \in [-1, 3]$ — (i)



& $x^2 - 2x - 2 \geq 0$

↳ $D = 4 + 8 = 12$

∴ $x = \frac{2 \pm 2\sqrt{3}}{2} = 1 \pm \sqrt{3}$

$(x - (1 + \sqrt{3})) \cdot (x - (1 - \sqrt{3})) \geq 0$

∴ $x \in (-\infty, 1 - \sqrt{3}] \cup [1 + \sqrt{3}, \infty)$



∴ $x \in [-1, 1 - \sqrt{3}, 1 + \sqrt{3}, 3]$

(Ans)

→ **(b)** $\log_5(x^2 - 11x + 43) < 2$

⇒ $x^2 - 11x + 43 < 25$

⇒ $x^2 - 11x + 18 < 0$

⇒ $(x-9)(x-2) < 0$

⇒ $x \in (2, 9)$

& $x^2 - 11x + 43 > 0$

↳ $D = 121 - 172 < 0$

always true

$x \in \mathbb{R}$

∴ $x \in 2, 9$



$$2 - \log_2(x^2 + 3x) \geq 0$$

$$\Rightarrow \log_2(x^2 + 3x) \leq 2$$

$$\Rightarrow x^2 + 3x \leq 4$$

$$\Rightarrow x^2 + 3x - 4 \leq 0$$

$$\Rightarrow (x+4)(x-1) \leq 0$$

$$\therefore x \in [-4, 1] \quad \text{--- (1)}$$

$$x^2 + 3x > 0$$

$$\Rightarrow x(x+3) > 0$$

$$\Rightarrow x \in (-\infty, -3) \cup (0, \infty)$$

$$\text{Ans. } x \in [-4, -3) \cup (0, 1]$$

Ans.

$$\log_{1.5} \frac{2x-8}{x-2} < 0$$

$$\Rightarrow \frac{2x-8}{x-2} - 1 < 0$$

$$\Rightarrow \frac{x-6}{x-2} < 0$$

$$\Downarrow x \in (2, 6)$$

&

$$\frac{2x-8}{x-2} > 0$$

$$\Rightarrow \frac{x-4}{x-2} > 0 \Rightarrow x \in (-\infty, 2) \cup (4, \infty)$$

$$\text{Ans. } x \in (4, 6)$$



→ (e) $\log_3 \frac{1+2x}{1+x} < 1$

⇒ $\log_3 \frac{1+2x}{1+x} < 3$

⇒ $\frac{1+2x-3-3x}{1+x} < 0$

⇒ $\frac{x+2}{1+x} > 0$

∴ $x \in (-\infty, -2) \cup (-1, \infty)$ — (i)

&

$\frac{1+2x}{1+x} > 0$

∴ $x \in (-\infty, -1) \cup (-\frac{1}{2}, \infty)$ — (ii)

∴

∴ $x \in (-\infty, -2) \cup (-\frac{1}{2}, \infty)$ Ans.

→ (f) $\log_4 \frac{3x+2}{x} \leq 0.5$

⇒ $\frac{3x+2}{x} \leq \sqrt{4} = 2$ & $\frac{3x+2}{x} > 0$

⇒ $\frac{3x+2-2x}{x} \leq 0$

⇒ $\frac{x+2}{x} \leq 0$

∴ $x \in [-2, 0)$

&

∴ $x \in (-\infty, -\frac{2}{3}) \cup (0, \infty)$

∴

∴ $x \in [-2, -\frac{2}{3})$ Ans.




→ Q1) $\log_4 \frac{3x+2}{x} \cdot \log_2 \left(\frac{x^2-4x+2}{x+1} \right) \leq 1$

$$\Rightarrow \frac{x^2-4x+2}{x+1} \leq 2$$

$$\Rightarrow \frac{x^2-4x+2-2x-2}{x+1} \leq 0$$

$$\Rightarrow \frac{x^2-6x}{x+1} \leq 0$$



$$\Rightarrow \frac{x(x-6)}{x+1} \leq 0$$



& $\frac{x^2-4x+2}{x+1} > 0$

$D = 16 - 8 = 8$
 $\therefore x = 2 \pm \sqrt{2}$

$$\Rightarrow \frac{(x - (2 + \sqrt{2}))(x - (2 - \sqrt{2}))}{x+1} > 0$$



Ans ⇒ $\therefore x \in [0, 2 - \sqrt{2}) \cup (2 + \sqrt{2}, 6]$



Solution to Previous KTKs

ATDB.uno

QUESTION [MHT CET 2023 (9 May)]

(KTK 01)



If $\log_2 x + \log_4 x + \log_8 x + \log_{16} x = \frac{25}{36}$ and $x = 2^k$ then k is

A 1

B $\frac{1}{2}$

C $\frac{1}{3}$

D $\frac{1}{8}$

ATDB.uno

Ans. C

KTK-1.

Q. If $\log_2 x + \log_4 x + \log_8 x + \log_{16} x = \frac{25}{36}$ and

$x = 2^k$ then:

$$\Rightarrow \log_2 x + \frac{1}{2} \log_2 x + \frac{1}{3} \log_2 x + \frac{1}{4} \log_2 x = \frac{25}{36}$$

let, $\log_2 x = t$

krish keshri
jharkhand

$$\Rightarrow t + \frac{t}{2} + \frac{t}{3} + \frac{t}{4} = \frac{25}{36} \Rightarrow \frac{12t + 6t + 4t + 3t}{12} = \frac{25}{36}$$

$$\Rightarrow \frac{25t}{12} = \frac{25}{36} \Rightarrow t = \frac{12}{36} = \frac{1}{3}$$

$$\# \log_2 x = \frac{1}{3}$$

$$\Rightarrow x = (2)^{1/3} \Rightarrow x = 2^k = 2^{1/3}$$

* $k = 1/3$ Ans.

$$t = \frac{1}{3}$$

QUESTION [BITSAT 2021]

(KTK 02)



If $\log_7 5 = a$, $\log_5 3 = b$ and $\log_3 2 = c$, then the logarithm of the number 70 to the base 225 is

A $\frac{1 - a + abc}{2a(1 + b)}$

B $\frac{1 - a - abc}{2a(1 + b)}$

C $\frac{1 + a - abc}{2a(1 + b)}$

D $\frac{1 + a + abc}{2a(1 + b)}$

ATDB.uno

Ans. D



the logarithm of the number 70 to the base

225 is : $\neq \log_7 5$, $\log_5 3$, $\log_3 2 = \log_7 2$
 $\hookrightarrow (abc)$

$\Rightarrow \log_{225} 70 \Rightarrow \frac{\log_2 70}{\log_2 225}$ $\neq \log_5 3$, $\log_3 2 = \log_5 2$
 $\hookrightarrow (bc)$

$\Rightarrow \frac{\log_2 (2 \times 5 \times 7)}{\log_2 (3 \times 3 \times 5 \times 5)}$ $\Rightarrow \frac{(bc)(abc) + (abc) + (bc)}{(bc)(abc)}$

$\Rightarrow \frac{\log_2 2 + \log_2 5 + \log_2 7}{\log_2 3^2 + \log_2 5^2}$ $\Rightarrow \frac{bc + 2c}{(c)(bc)}$

$\Rightarrow \frac{1 + \frac{1}{\log_5 2} + \frac{1}{\log_7 2}}{2 \left(\frac{1}{\log_3 2} + \frac{1}{\log_5 2} \right)}$ $\Rightarrow \frac{(bc) [(abc) + a + 1]}{(bc)(abc)} \times \frac{(c)(bc)}{2c(1+b)}$

$\Rightarrow \frac{1 + \frac{1}{bc} + \frac{1}{abc}}{2 \left(\frac{1}{c} + \frac{1}{bc} \right)}$ $\Rightarrow \frac{1 + a + abc}{2a(1+b)}$ Ans.

krish keshri
iharkhand

KTK-2) If $\log_7^5 = a$, $\log_5^3 = b$ and $\log_3^2 = c$, then the logarithm of the number 70 to the base 225 is:

$$\log_7^5 \times \log_5^3 \times \log_3^2 = \log_7^2 = abc$$

$$\log_{225}^{70} = \frac{\log_7^{70}}{\log_7^{225}} = \frac{\log_7^7 + \log_7^{10}}{\log_7^{25} + \log_7^9}$$

$$\log_7^5 \times \log_5^3 = \log_7^3 = ab$$

$$= \frac{1 + \log_7^2 + \log_7^5}{2 \log_7^5 + 2 \log_7^3}$$

$$= \frac{1 + abc + a}{2(\log_7^5 + \log_7^3)}$$

**Kriti Mathur
Raj.**

$$= \frac{1 + abc + a}{2(a + ab)}$$

$$\log_{225}^{70} = \frac{1 + a + abc}{2a(1 + b)} \quad \textcircled{D} \quad \underline{\underline{\text{Ans.}}}$$



QUESTION [BITSAT 2020]

(KTK 03)



If $\log_5 \frac{(a + b)}{3} = \frac{\log_5 a + \log_5 b}{2}$, then $\frac{a^4 + b^4}{a^2 b^2}$ is equal to

A 50

B 47

C 44

D 53

ATDB.uno

Ans. B



KTK-03.

Q If $\log_5 \frac{(a+b)}{3} = \frac{\log_5 a + \log_5 b}{2}$, then $\frac{a^4 + b^4}{a^2 b^2}$ is equal to:

$\Rightarrow 2 \log_5 \frac{(a+b)}{3} = \log_5 (a \cdot b)$

$\Rightarrow \log_5 \left(\frac{(a+b)}{3}\right)^2 = \log_5 (a \cdot b)$

$\Rightarrow a^2 + b^2 + 2ab = 3ab$

$\Rightarrow \boxed{a^2 + b^2 = ab}$
SBS.

$\Rightarrow a^4 + b^4 + 2a^2 b^2 = 49 a^2 b^2$

$\Rightarrow a^4 + b^4 = 47 a^2 b^2$

$\Rightarrow \boxed{\frac{a^4 + b^4}{a^2 b^2} = 47}$ Ans.

krish keshri
jharkhand



KTK-3) If $\log_5 \left(\frac{a+b}{3} \right) = \frac{\log_5 a + \log_5 b}{2}$, then $\frac{a^4 + b^4}{a^2 b^2}$ is equal

to

A.) 50

B.) 47

C.) 44

D.) 53

$$\log_5 \left(\frac{a+b}{3} \right) = \frac{\log_5 a + \log_5 b}{2}$$

$$\log_5 \left(\frac{a+b}{3} \right) = \frac{\log_5 ab}{2}$$

$$2 \log_5 \left(\frac{a+b}{3} \right) = \log_5 ab \Rightarrow \log_5 \left(\frac{a+b}{3} \right)^2 = \log_5 ab$$

**Kriti Mathur
Raj.**

ATDB.uno

$$\Rightarrow \frac{(a+b)^2}{9} = ab$$

$$\Rightarrow a^2 + b^2 + 2ab - 9ab = 0$$

$$\Rightarrow a^2 + b^2 = 7ab$$

$$\text{Now, } \frac{a^4 + b^4}{a^2 b^2} = \frac{(a^2 + b^2)^2 - 2a^2 b^2}{a^2 b^2}$$

$$= \frac{49a^2 b^2 - 2a^2 b^2}{a^2 b^2} = \frac{47a^2 b^2}{a^2 b^2}$$

$$= 47 \text{ (B) Ans.}$$

QUESTION [WB JEE 2024]

(KTK 04)



If $(x^2 \log_x 27) \cdot \log_9 x = x + 4$ then the value of x is

A 2

B $-\frac{4}{3}$

C -2

D $\frac{4}{3}$

ATDB.uno

Ans. A

KTK-04.

If $(x^2 \log_x 27) \cdot \log_9 x = x+4$ then the value of x is :

$$\Rightarrow x^2 \cdot \frac{\log_3 3^3}{\log_3 x} \cdot \frac{1}{2} \log_3 x = x+4$$

**krish keshri
jharkhand**

$$\Rightarrow \frac{3x^2}{2} = x+4 \Rightarrow 3x^2 - 2x - 8 = 0$$

$$\Rightarrow 3x^2 - 6x + 4x - 8 = 0$$

$$\Rightarrow 3x(x-2) + 4(x-2) = 0$$

$$\Rightarrow (x-2)(3x+4) = 0$$

$$\Rightarrow x = 2, \frac{-4}{3} \Rightarrow \boxed{x=2} \text{ Ans.}$$

↪ REJECT





KTK-40) If $(x^2 \log_a 27) \cdot \log_a x = x+4$ then the value of x is

A.) 2 B.) $-\frac{4}{3}$ C.) -2 D.) $\frac{4}{3}$

$$\Rightarrow x^2 \log_a 27 \cdot \log_a x = x+4$$

$$\Rightarrow \frac{x^2 \log_a 27}{\log_a x} = x+4$$

$$\Rightarrow \frac{x^2 \log_a 3^3}{\log_a 3^2} = x+4$$

$$\Rightarrow \frac{x^2 \times 3 \log_a 3}{2 \log_a 3} = x+4$$

$$\Rightarrow 3x^2 = 2x + 8$$

$$3x^2 - 2x - 8 = 0$$

$$3x^2 - 6x + 4x - 8 = 0$$

$$\Rightarrow 3x(x-2) + 4(x-2) = 0$$

$$(x-2)(3x+4) = 0$$

$$\textcircled{A} [x=2] \quad x = -\frac{4}{3} \times$$

Ans.

**Kriti Mathur
Raj.**

ATDB.uno

QUESTION [WB JEE 2020]

(KTK 05)



If $2 \log(x + 1) - \log(x^2 - 1) = \log 2$, then $x =$

- A** only 3
- B** -1 and 3
- C** only -1
- D** 1 and 3

ATDB.uno

Ans. A



KTK-05.

If $2 \log (x+1) - \log (x^2-1) = \log 2$, then $x =$.

$$\Rightarrow 2 \log \frac{(x+1)}{(x^2-1)} = \log 2$$

$$\Rightarrow \frac{(x+1)^2}{(x+1)(x-1)} = \log 2$$

$$\Rightarrow x+1 = 2x-2$$

$$\Rightarrow \boxed{x=3} \text{ Ans.}$$

krish keshri
jharkhand



KTK-5.) If $2 \log(x+1) - \log(x^2-1) = \log 2$, then $x =$

$$\log(x+1)^2 - \log(x^2-1) = \log 2$$

$$\log\left(\frac{(x+1)^2}{x^2-1}\right) = \log 2$$

$$\frac{(x+1)^2}{x^2-1} = 2$$

$$\frac{(x+1)^2}{(x+1)(x-1)} = 2$$

$$\frac{(x+1)}{(x-1)} = 2 ; x \neq -1$$

$$x+1 = 2x-2$$

$$0 = 2x - x - 2 - 1$$

$$0 = x - 3 \Rightarrow \underline{\underline{x = 3}} \text{ Ans.}$$

Kriti Mathur
Raj.

ATDB.uno

QUESTION [WB JEE 2018]

(KTK 06)



If $x + \log_{10}(1 + 2^x) = x \log_{10} 5 + \log_{10} 6$, then the value of x is

A $\frac{1}{2}$

B $\frac{1}{3}$

C 1

D 2

ATDB.uno

Ans. C



KTK-OG

Q. If $x + \log_{10}(1+2^x) = x \log_{10} 5 + \log_{10} 6$,
then the value of x is.

$$\Rightarrow x \log_{10} 10 + \log_{10}(1+2^x) = \log_{10} 5^x + \log_{10} 6$$

$$\Rightarrow \log_{10} [10^x (1+2^x)] = \log_{10} (5^x \cdot 6)$$

$$\Rightarrow 10^x \cdot (1+2^x) = 5^x \cdot 6$$

$$\Rightarrow 2^x (1+2^x) = 6$$

ATDB.uno

let, $2^x = t$

krish keshri
jharkhand

$$\Rightarrow t(1+t) = 6$$

$$\Rightarrow t^2 + t - 6 = 0 \Rightarrow (t+3)(t-2) = 0$$

$$\Rightarrow t = -3, 2$$

$$\Rightarrow 2^x = -3 \quad | \quad \Rightarrow 2^x = 2$$

$$\underline{\text{NOT POSSIBLE}} \quad | \quad \Rightarrow \boxed{x=1} \quad \underline{\text{Ans}}$$



KTK-6) If $x + \log_{10} (1+2^x) = x \log_{10} 5 + \log_{10} 6$, then the value of x is

Ans. $x + \log_{10} (1+2^x) = \log_{10} 5^x + \log_{10} 6$

$x + \log_{10} (1+2^x) = \log_{10} 5^x \cdot 6$

$\Rightarrow x = \log_{10} 5^x \cdot 6 - \log_{10} (1+2^x)$

**Kriti Mathur
Raj.**

$\Rightarrow \log_{10} \left(\frac{5^x \cdot 6}{2^x + 1} \right) = x$

$\frac{5^x \cdot 6}{2^x + 1} = 10^x$

Now, by hit & trial

at $x = 1 \Rightarrow \frac{5 \cdot 6}{2+1} = \frac{30}{3} \Rightarrow 10 = 10$ **(C) Ans.**
[$x=1$]

QUESTION [WB JEE 2019]

(KTK 07)



If $\log_2 6 + \frac{1}{2x} = \log_2 (2^{\frac{1}{x}} + 8)$, then the value of x are

A $\frac{1}{4}, \frac{1}{3}$

B $\frac{1}{4}, \frac{1}{2}$

C $-\frac{1}{4}, \frac{1}{2}$

D $\frac{1}{3}, -\frac{1}{2}$

ATDB.uno

Ans. B



KTK-07.

Q. If $\log_2 6 + \frac{1}{2x} = \log_2 (2^{1/x} + 8)$, then the

value of x are : write: $\frac{1}{2x} = \log_2 (2^{1/2x})$

$$\Rightarrow \log_2 (6 \cdot 2^{1/2x}) = \log_2 (2^{1/x} + 8)$$

$$\Rightarrow 6 \cdot 2^{1/2x} = 2^{1/x} + 8 \quad \text{let; } 2^{1/x} = t$$

$$\Rightarrow 6 t^{1/2} = t + 8$$

SBS

ATDB.uno

krish keshri
jharkhand

$$\Rightarrow 36 t \quad t^2 + 16t + 64$$

$$\Rightarrow t^2 - 20t + 64 = 0$$

$$\Rightarrow t^2 - 16t - 4t + 64 = 0$$

$$\Rightarrow t(t-16) - 4(t-16) = 0$$

$$\Rightarrow (t-16)(t-4) = 0$$

$$\Rightarrow t = 16, 4.$$

$$\Rightarrow 2^{1/x} = 16 \quad | \quad \Rightarrow 2^{1/x} = 4$$

$$\Rightarrow \frac{1}{x} = 4 \quad | \quad \Rightarrow \frac{1}{x} = 2$$

$$\Rightarrow \boxed{x = \frac{1}{4}} \quad | \quad \Rightarrow \boxed{x = \frac{1}{2}}$$

Ans.



KTK-7.) If $\log_2 6 + \frac{1}{2x} = \log_2 (2^{\frac{1}{2}} + 8)$, then the value of x are.

A.) $\frac{1}{4}, \frac{1}{3}$

~~B.) $\frac{1}{4}, \frac{1}{2}$~~

C.) $-\frac{1}{4}, \frac{1}{2}$

D.) $\frac{1}{3}, -\frac{1}{2}$

$$\log_2 6 + \frac{1}{2x} = \log_2 (2^{\frac{1}{2}} + 8)$$

$$\log_2 6 + \log_2 2^{\frac{1}{2x}} = \log_2 (2^{\frac{1}{2}} + 8)$$

$$\log_2 (6 \times 2^{\frac{1}{2x}}) = \log_2 (2^{\frac{1}{2}} + 8)$$

$$6 \times 2^{\frac{1}{2x}} = 2^{\frac{1}{2}} + 8$$

$$\text{Let } 2^{\frac{1}{2x}} = t, \quad t^2 = 2^{\frac{1}{x}}$$

$$6t = t^2 + 8$$

$$t^2 - 6t + 8 = 0$$

$$t^2 - 4t - 2t + 8 = 0$$

$$t(t-4) - 2(t-4) = 0$$

$$t = 2$$

$$2^{\frac{1}{2x}} = 2$$

$$\frac{1}{2x} = 1$$

$$2x = 1$$

$$\left[x = \frac{1}{2} \right] \underline{\underline{\text{Ans.}}}$$

$$t = 4$$

$$2^{\frac{1}{2x}} = 2^2$$

$$\frac{1}{2x} = 2$$

$$\left[x = \frac{1}{4} \right] \underline{\underline{\text{Ans.}}}$$

$$\textcircled{B} \underline{\underline{\text{Ans.}}}$$

**Kriti Mathur
Raj.**

ATDB.uno

QUESTION [WB JEE 2017]

(KTK 08)



If $(\log_5 x)(\log_x 3x)(\log_{3x} y) = \log_x x^3$, then y equals

A 125

B 25

C 513

D 243

ATDB.uno

Ans. A



KTK-08.

If $(\log_5 x) (\log_x 3x) (\log_{3x} y) = \log_x x^3$ then

y equals :

$$\Rightarrow \log_5 x \cdot \log_x 3x \cdot \log_{3x} y = 3$$

$$\Rightarrow \log_5 y = 3$$

$$\Rightarrow y = 5^3 \Rightarrow \boxed{y = 125} \text{ Ans.}$$

krish keshri
jharkhand

QUESTION [COMEDK 2023]

(KTK 09)



The value of $a^{\log_b c} - c^{\log_b a}$, where $a, b, c > 0$ but $a, b, c \neq 1$, is

A a

B b

C c

D 0

ATDB.uno

Ans. D



KTK-09.

The value of $a^{\log_b c} - c^{\log_b a}$, where $a, b, c > 0$

but $a, b, c \neq 1$, is:

$$\Rightarrow a^{\log_b c} - a^{\log_b c} \quad \text{interchange.}$$

$$\Rightarrow 0 \quad \underline{\text{Ans.}}$$

QUESTION [COMEDK 2022]

(KTK 10)



The value of $3^{\log_4 5} - 5^{\log_4 3}$ is

A 0

B 1

C 2

D 4

ATDB.uno

Ans. A



KTK-10, The value of $3^{\log_4 5} - 5^{\log_4 3}$ is :

$$\Rightarrow 3^{\log_4 5} - 3^{\log_4 5} \leftarrow \text{interchange.}$$

$$\Rightarrow 0 \text{ Ans.}$$

QUESTION [COMEDK 2021]

(KTK 11)



$8^3 \log_8 5$ is equal to

A $\log_8 25$

B 120

C 125

D $\log_8 15$

ATDB.uno

Ans. C



KTK-11.

$8^3 \log_8 5$ is equal to :

$$\Rightarrow 8^{\log_8 5^3}$$

$$\Rightarrow 5^3 = 125 \text{ Ans.}$$

krish keshri
jharkhand

QUESTION [COMEDK 2020]

(KTK 12)



$7^{2 \log_7 5}$ is equal to

- A** 5
- B** $\log_7 35$
- C** $\log_7 25$
- D** 25

ATDB.uno

Ans. D



KTK-12.

$7^2 \log_7 5$ is equal to:

$$\Rightarrow 7 \log_7 5^2$$

$$\Rightarrow 5^2 = 25 \text{ Ans.}$$

QUESTION

(KTK 13)



Find the exhaustive solutions set of $\frac{(x^2-9)^{101}(x^2+6)(x^2-4)^{100}}{(x^2-5x+6)^{13}(x^2-16)^{16}} > 0$.

ATDB.uno

Ans. $(-\infty, -3) \cup (2, \infty) - \{\pm 4, 3\}$



KTK-13) Find the exhaustive solutions set of $\frac{(x^2+9)^{101} (x^2+6) (x^2-4)^{100}}{(x^2-5x+6)^{13} (x^2-16)^{16}} > 0$.

$$\Rightarrow \frac{(x+3)^{101} (x-3)^{101} (x^2+6)}{(x^2-5x+6)^{13}} > 0, \quad x \neq \pm 2, \pm 4$$


$$\Rightarrow \frac{(x+3)^{101} (x-3)^{101} (x^2+6)}{(x-2)^3 (x-3)^3} > 0, \quad x \neq \pm 2, \pm 4$$

$$\Rightarrow \frac{(x+3)^{101} (x-3)^{89} (x^2+6)^{atongh+ve}}{(x-2)^3} > 0, \quad x \neq \pm 2, \pm 4, 3$$

ATDB.uno

Kriti Mathur
Raj.

$$\Rightarrow \frac{(x+3)^{101}}{(x-2)^3} > 0$$



$$x \in (-\infty, -3) \cup (2, \infty) - \{\pm 4, 3\} \underline{\underline{\text{Ans.}}}$$

QUESTION

(KTK 14)



Find the exhaustive solutions set of $\frac{(x-4)^{30}(x^2-9)^9(x^2-3x+2)^{17}(3x^2+10)^{10}}{(x^2-5x+6)^{52}(x^2-25)^{60}(x^2+10)^{11}} \leq 0$.

ATDB.uno

Ans. $[-3, 1] \cup (2, 3)$



KTK-14) Find the exhaustive solutions set of

$$\frac{(x-4)^{30} (x^2-9)^9 (x^2-3x+2)^{17} (3x^2+10)^{10}}{(x^2-5x+6)^{52} (x^2-25)^{60} (x^2+10)^{11}} \leq 0$$


$$\Rightarrow \frac{(x+3)^9 (x-3)^9 (x^2-3x+2)^{17}}{(x^2-5x+6)^{52} (x^2+10)^{11}} \leq 0; x \neq \pm 5, x=4$$

$$\Rightarrow \frac{(x+3)^9 (x-3)^9 (x-2)^{17} (x-1)^{17}}{(x-2)^{52} (x-3)^{42}} \leq 0; x \neq \pm 5$$

$$\Rightarrow \frac{(x+3)^9 (x-1)^{17}}{(x-2)^{35} (x-3)^{42}} \leq 0; x \neq \pm 5$$

Kriti Mathur

Raj.



$$x \in [-3, 1] \cup (2, 3) \cup \{4\}$$

QUESTION

(KTK 15)



Solve in real numbers the equation $\sqrt{x} + \sqrt{y} + 2\sqrt{z-2} + \sqrt{u} + \sqrt{v} = x + y + z + u + v$.

$$0 = x - 2 \cdot \sqrt{x} \cdot \frac{1}{2} + \frac{1}{4} + y - 2\sqrt{y} \cdot \frac{1}{2} + \frac{1}{4} + u - 2\sqrt{u} \cdot \frac{1}{2} + \frac{1}{4} + v - 2\sqrt{v} \cdot \frac{1}{2} + \frac{1}{4} + \sqrt{z-2}^2 - 2\sqrt{z-2} + 1 + z - \frac{1}{4} - \frac{1}{4} - \frac{1}{4} - \frac{1}{4} - 1$$

$$0 = (\sqrt{x} - 1/2)^2 + (\sqrt{y} - 1/2)^2 + (\sqrt{u} - 1/2)^2 + (\sqrt{v} - 1/2)^2 + (\sqrt{z-2} - 1)^2 = 0$$

ATDB.uno

$$x = 1/4 = y = u = v$$

$$z = 3$$

Ans. $x = y = u = v = 1/4, z = 3$

KTK-15

Solve in real no.s, the eqn

$$\sqrt{x} + \sqrt{y} + 2\sqrt{z-2} + \sqrt{u} + \sqrt{v} = x + y + z + u + v.$$

KTK 15

by Reed

from WB

Soln

$$x + y + z + u + v = \sqrt{x} + \sqrt{y} + 2\sqrt{z-2} + \sqrt{u} + \sqrt{v}$$

$$\Rightarrow x - \sqrt{x} + y - \sqrt{y} + z - 2\sqrt{z-2} + u - \sqrt{u} + v - \sqrt{v} = 0$$

$$\Rightarrow x - 2 \cdot \sqrt{x} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2 + y - 2\sqrt{y} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2 + (z-2) - 2\sqrt{z-2} \cdot 1 + 1^2$$

$$+ u - 2\sqrt{u} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2 + v - 2\sqrt{v} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2$$

$$= \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 + 1^2 + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2$$

$$\Rightarrow \left(\sqrt{x} - \frac{1}{2}\right)^2 + \left(\sqrt{y} - \frac{1}{2}\right)^2 + \left(\sqrt{z-2} - 1\right)^2 + \left(\sqrt{u} - \frac{1}{2}\right)^2 + \left(\sqrt{v} - \frac{1}{2}\right)^2$$

$$= \frac{1}{4} + \frac{1}{4} + 1 + \frac{1}{4} + \frac{1}{4} - 2 = 2 - 2 = 0.$$

$$\Rightarrow \left. \begin{array}{l} \sqrt{x} = \frac{1}{2} \\ x = \frac{1}{4} \end{array} \right| \left. \begin{array}{l} \sqrt{y} = \frac{1}{2} \\ y = \frac{1}{4} \end{array} \right| \left. \begin{array}{l} \sqrt{z-2} = 1 \\ z-2 = 1 \\ z = 3 \end{array} \right| \left. \begin{array}{l} \sqrt{u} = \frac{1}{2} \\ u = \frac{1}{4} \end{array} \right| \left. \begin{array}{l} \sqrt{v} = \frac{1}{2} \\ v = \frac{1}{4} \end{array} \right|$$

$$\therefore \boxed{x = y = u = v = \frac{1}{4}} \text{ \& \ } \boxed{z = 3}$$



QUESTION**(KTK 16)**

Find all pair of positive integer (m, h) that satisfy $mn + 3m - 8n = 59$.

ATDB.uno**Ans. 3**



KTK-16) Find all pair of positive integers (m, n) that satisfy

$$mn + 3m - 8n = 59$$

Ans. $m(n+3) - 8n = 24 + 35$

$$m(n+3) - 8n - 24 = 35$$

$$m(n+3) - 8(n+3) = 35$$

$$(m-8)(n+3) = 35$$

59

24

35

$(1 \times 35, 5 \times 7,$

$7 \times 5, 35 \times 1$

Kriti Mathur
Raj.

ATDB.uno

$m-8$

$n+3$

(m, n)

1

35

$(9, 32)$

5

7

$(13, 4)$

7

5

$(15, 2)$

35

1

$(43, -2) \times$

3 pairs



KTK-161 Find all pair of +ve integer (m, n) that
 $mn + 3m - 8n = 59.$

Soln $mn + 3m - 8n = 59$

a. $m(n+3) - 8n - 24 = 59 - 24$

a. $m(n+3) - 8(n+3) = 35$

a. $(m-8)(n+3) = 35.$

now, $35 = 5 \times 7$

$35 = 1 \times 35$

✓

$35 = (-5) \times (-7)$

$35 = (-1) \times (-35)$

ATDB.uno

no need since $m, n \in \mathbb{I}^+$

$m-8 =$	5	7	1	35
$n+3 =$	7	5	35	1
$\Rightarrow m =$	13	15	9	43
$\& n =$	4	2	32	-2

✓ ✓ ✓ ✓
 (∵ $m, n \in \mathbb{I}^+$)

∴ all pair of positive integers $(m, n) \in (13, 4), (15, 2), (9, 32)$
 (Ans.)

∴ no. of pairs of (m, n) is = 3.

QUESTION

(KTK 17)



The least value of the expression $(x + y)(y + z)$ where given that $x, y, z > 0$ and $xyz(x + y + z) = 1$

ATDB.uno

Ans. 2

KTK-170) The least value of the expression $(x+y)(y+z)$ where given that $x, y, z > 0$ and $xyz(x+y+z) = 1$

Ans. $x+y+z = \frac{1}{xyz}$

**Kriti Mathur
Raj.**

$$(x+y)(y+z)$$

$$xy + xz + y^2 + yz$$

$$= y(x+y+z) + xz$$

ATDB.uno

$$\Rightarrow y \left(\frac{1}{xyz} \right) + xz$$

$$\Rightarrow \frac{1}{xz} + xz$$

$$\text{As } \frac{1}{xz} + xz \geq 2$$

Least value = 2 Ans.



Date: / /
Page: /



KTK-17! The least value of the expression
 $(x+y)(y+z)$ where given that $x, y, z > 0$.

and $xyz(x+y+z) = 1$.

Soln

$$E = (x+y)(y+z)$$

$$\Rightarrow E = xy + xz + yz + yz$$

$$\Rightarrow E = y(x+y+z) + yz$$

$$\Rightarrow E = \frac{y}{xyz} + xz \quad \neq y \neq 0$$

$$\Rightarrow E = \frac{1}{xz} + xz$$

$$\therefore E_{\min} \Big|_{\text{at } x=z=1} = 2$$

$$\therefore \boxed{E_{\min} = 2} \text{ Ans.}$$

$$\left. \begin{aligned} xyz(x+y+z) &= 1 \\ \Rightarrow (x+y+z) &= \frac{1}{xyz} \end{aligned} \right\}$$

KTK 17

by Reed

from WB



* Read class Theory

* Retry class Questions In rough COPY

* TAH

* BPP

* KTK

* DPP / Module.

ATDB.uno



THANK ATDB.uno YOU