


## CHAPTER

## 4



# Trigonometric Ratios and Identities

## 1. Trigonometric Ratio – Basic Terminology



The six trigonometric ratios of  $\theta$  are defined as follows:

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}, \quad \cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}, \quad \tan \theta = \frac{\text{opposite}}{\text{adjacent}}$$

$$\cot \theta = \frac{\text{adjacent}}{\text{opposite}}, \quad \operatorname{cosec} \theta = \frac{\text{hypotenuse}}{\text{opposite}}, \quad \sec \theta = \frac{\text{hypotenuse}}{\text{adjacent}}$$

## 2. Trigonometric identities:

- (i)  $\sin^2 \theta + \cos^2 \theta = 1$
- (ii)  $\sec^2 \theta - \tan^2 \theta = 1$
- (iii)  $\operatorname{cosec}^2 \theta - \cot^2 \theta = 1$

## 3. Allied angles:

Two angles are said to be allied when their sum or difference is either zero or a multiple of  $\frac{\pi}{2}$ , two angles  $x, y$  are allied angles iff  $|x \pm y| = 0$  or  $\frac{n\pi}{2}, n \in \mathbb{N}$ .

$q \rightarrow$	$\frac{\pi}{2} - \theta$	$\frac{\pi}{2} + \theta$	$\pi - \theta$	$\pi + \theta$	$\frac{3\pi}{2} - \theta$	$\frac{3\pi}{2} + \theta$	$2\pi - \theta$	$2\pi + \theta$	$-\theta$
sin	$\cos \theta$	$\cos \theta$	$\sin \theta$	$-\sin \theta$	$-\cos \theta$	$-\cos \theta$	$-\sin \theta$	$\sin \theta$	$-\sin \theta$
cos	$\sin \theta$	$-\sin \theta$	$-\cos \theta$	$-\cos \theta$	$-\sin \theta$	$\sin \theta$	$\cos \theta$	$\cos \theta$	$\cos \theta$
tan	$\cot \theta$	$-\cot \theta$	$-\tan \theta$	$\tan \theta$	$\cot \theta$	$-\cot \theta$	$-\tan \theta$	$\tan \theta$	$-\tan \theta$
cot	$\tan \theta$	$-\tan \theta$	$-\cot \theta$	$\cot \theta$	$\tan \theta$	$-\tan \theta$	$-\cot \theta$	$\cot \theta$	$-\cot \theta$
sec	$\operatorname{cosec} \theta$	$-\operatorname{cosec} \theta$	$-\sec \theta$	$-\sec \theta$	$-\operatorname{cosec} \theta$	$\operatorname{cosec} \theta$	$\sec \theta$	$\sec \theta$	$\sec \theta$
cosec	$\sec \theta$	$\sec \theta$	$\operatorname{cosec} \theta$	$-\operatorname{cosec} \theta$	$-\sec \theta$	$-\sec \theta$	$-\operatorname{cosec} \theta$	$\operatorname{cosec} \theta$	$-\operatorname{cosec} \theta$

## 4. Sum & Difference Formula

- (i)  $\sin(A + B) = \sin A \cos B + \cos A \sin B$
- (ii)  $\sin(A - B) = \sin A \cos B - \cos A \sin B$
- (iii)  $\cos(A + B) = \cos A \cos B - \sin A \sin B$
- (iv)  $\cos(A - B) = \cos A \cos B + \sin A \sin B$
- (v)  $\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$
- (vi)  $\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$
- (vii)  $\cot(A + B) = \frac{\cot B \cot A - 1}{\cot B + \cot A}$
- (viii)  $\cot(A - B) = \frac{\cot B \cot A + 1}{\cot B - \cot A}$

## 5. Product to sum

- (i)  $2 \sin A \cos B = \sin(A + B) + \sin(A - B)$
- (ii)  $2 \cos A \sin B = \sin(A + B) - \sin(A - B)$
- (iii)  $2 \cos A \cos B = \cos(A + B) + \cos(A - B)$
- (iv)  $2 \sin A \sin B = \cos(A - B) - \cos(A + B)$

## 6. Trigonometric transformations:

- (i)  $\sin C + \sin D = 2 \sin\left(\frac{C+D}{2}\right) \cos\left(\frac{C-D}{2}\right)$
- (ii)  $\sin C - \sin D = 2 \cos\left(\frac{C+D}{2}\right) \sin\left(\frac{C-D}{2}\right)$
- (iii)  $\cos C + \cos D = 2 \cos\left(\frac{C+D}{2}\right) \cos\left(\frac{C-D}{2}\right)$

$$(iv) \cos C - \cos D = -2 \sin\left(\frac{C+D}{2}\right) \sin\left(\frac{C-D}{2}\right)$$

$$= 2 \sin\left(\frac{C+D}{2}\right) \sin\left(\frac{D-C}{2}\right)$$

Some useful Formulae:

- (i)  $\sin(A+B) \sin(A-B) = \sin^2 A - \sin^2 B = \cos^2 B - \cos^2 A$
- (ii)  $\cos(A+B) \cos(A-B) = \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A$
- (iii)  $\tan(A+B) \tan(A-B) = \frac{\tan^2 A - \tan^2 B}{1 - \tan^2 A \tan^2 B}$
- (iv)  $\cot(A+B) \cot(A-B) = \frac{\cot^2 B \cot^2 A - 1}{\cot^2 B - \cot^2 A}$

7. Some standard trigonometric values

Angle → Trigonometric Function ↓	15°	18°	22½°	36°
sin	$\frac{\sqrt{3}-1}{2\sqrt{2}}$	$\frac{\sqrt{5}-1}{4}$	$\frac{1}{2}\sqrt{2-\sqrt{2}}$	$\frac{\sqrt{10-2\sqrt{5}}}{4}$
cos	$\frac{\sqrt{3}+1}{2\sqrt{2}}$	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{1}{2}\sqrt{2+\sqrt{2}}$	$\frac{\sqrt{5}+1}{4}$
tan	$2-\sqrt{3}$	$\frac{\sqrt{25-10\sqrt{5}}}{5}$	$\sqrt{2}-1$	$\sqrt{5-2\sqrt{5}}$

8. Double Angle / Triple Angle

- (i)  $\sin 2A = 2 \sin A \cos A$
  - (ii)  $\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$
- Suppose that  $A$  is not an odd multiple of  $\frac{\pi}{4}$ . Then
- (iii)  $\sin 2A = \frac{2 \tan A}{1 + \tan^2 A}$
  - (iv)  $\cos 2A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$
  - (v)  $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$  (Here  $2A$  is also not an odd multiple of  $\frac{\pi}{2}$ )
  - (vi)  $\sin 3A = 3 \sin A - 4 \sin^3 A, \forall A \in R$
  - (vii)  $\cos 3A = 4 \cos^3 A - 3 \cos A, \forall A \in R$
  - (viii)  $\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$  ( $3A, A$  are not odd multiples of  $\frac{\pi}{2}$ )
  - (ix)  $\cot 3A = \frac{3 \cot A - \cot^3 A}{1 - 3 \cot^2 A}$  ( $3A, A$  are not multiples of  $\frac{\pi}{2}$ )

9. Half Angle

- (i)  $\sin \frac{A}{2} = 2 \sin \frac{A}{4} \cos \frac{A}{4}$
- (ii)  $\cos \frac{A}{2} = \cos^2 \frac{A}{4} - \sin^2 \frac{A}{4} = 2 \cos^2 \frac{A}{4} - 1 = 1 - 2 \sin^2 \frac{A}{4}$
- (iii)  $\tan \frac{A}{2} = \frac{2 \tan \frac{A}{4}}{1 - \tan^2 \frac{A}{4}}$  (where  $\frac{A}{2} \neq (2n+1)\frac{\pi}{2}, n \in Z$ )

10. Conditional Identities

In a Triangle If  $A + B + C = \pi$ , then

- (i)  $\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C$
- (ii)  $\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}$
- (iii)  $\cos 2A + \cos 2B + \cos 2C = -1 - 4 \cos A \cos B \cos C$
- (iv)  $\sin^2 A + \sin^2 B + \sin^2 C = 2(1 + 4 \cos A \cos B \cos C)$
- (v)  $\cos^2 A + \cos^2 B + \cos^2 C = (1 - 2 \cos A \cos B \cos C)$

11. Domains, Ranges and Periodicity of Trigonometric Functions:

T-Ratio	Domain	Range	Period
sin x	R	[-1, 1]	2π
cos x	R	[-1, 1]	2π
tan x	$R - \left\{ (2n+1)\frac{\pi}{2} \right\}; n \in I$	R	π
cot x	$R - \{n\pi: n \in I\}$	R	π
sec x	$R - \left\{ (2n+1)\frac{\pi}{2} \right\}; n \in I$	$(-\infty, -1] \cup [1, \infty)$	2π
cosec x	$R - \{n\pi: n \in I\}$	$(-\infty, -1] \cup [1, \infty)$	2π

12. Maxima-Minima

- (i)  $a \cos \theta + b \sin \theta$  will always lie in the interval  $[-\sqrt{a^2+b^2}, \sqrt{a^2+b^2}]$  i.e., the maximum and minimum values are  $\sqrt{a^2+b^2}, -\sqrt{a^2+b^2}$  respectively.
- (ii) Minimum value of  $a^2 \tan^2 \theta + b^2 \cot^2 \theta$  is  $2ab$  where  $a, b > 0$ .
- (iii)  $-\sqrt{a^2+b^2+2ab \cos(\alpha-\beta)} < a \cos(\alpha+\theta) + b \cos(\beta+\theta) \leq \sqrt{a^2+b^2+2ab \cos(\alpha-\beta)}$  where  $\alpha$  and  $\beta$  are known angles.



- (iv) If  $\alpha, \beta, \in \left(0, \frac{\pi}{2}\right)$  and  $\alpha + \beta = \sigma$  (constant) then
- (a) Maximum value of the expression  $\cos \alpha \cos \beta, \cos \alpha + \cos \beta, \sin \alpha \sin \beta$  or  $\sin \alpha + \sin \beta$  occurs when  $\alpha = \beta = \frac{\sigma}{2}$
- (b) Minimum value of  $\sec \alpha + \sec \beta, \tan \alpha + \tan \beta, \operatorname{cosec} \alpha + \operatorname{cosec} \beta$  occurs when  $\alpha = \beta = \frac{\sigma}{2}$
- (v) If  $A, B, C$  are the angles of a triangle then maximum value of  $\sin A + \sin B + \sin C$  and  $\sin A \sin B \sin C$  occurs when  $A = B = C = 60^\circ$
- (vi) In case a quadratic in  $\sin \theta$  &  $\cos \theta$  is given then the maximum or minimum values can be obtained by making perfect square.
13. Sum of Three or More Angles
- (i)  $\sin(A + B + C) = \sin A \cos B \cos C + \cos A \sin B \cos C + \cos A \cos B \sin C - \sin A \sin B \sin C = \sum \sin A \cos B \cos C - \prod \sin A$

$$(ii) \cos(A + B + C) = \cos A \cos B \cos C - \cos A \sin B \sin C - \sin A \cos B \sin C - \sin A \sin B \sin C = \prod \cos A - \sum \cos A \sin B \sin C$$

$$(iii) \tan(A + B + C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A} = \frac{\sum \tan A - \prod \tan A}{1 - \sum \tan A \tan B}$$

## 14. Summation of Series

$$\sin \alpha + \sin(\alpha + \beta) + \sin(\alpha + 2\beta) + \dots + \sin(\alpha + (n-1)\beta) = \frac{\sin\left(\alpha + \frac{n-1}{2}\beta\right) \sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}}$$

$$\cos \alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos(\alpha + (n-1)\beta) = \frac{\cos\left(\alpha + \frac{n-1}{2}\beta\right) \sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}}$$

# ATDB.uno